

Since 2013 indexed in



IBI Factor 2014-15 is 3.43



Since 2015 indexed in

EBSCO Academic Search Ultimate & One Belt, One Road Reference Source

SUMMARIES OF THE 12TH NATIONAL CONFERENCE OF THE ROMANIAN SOCIETY OF OBSTETRICS AND GYNECOLOGY
• Cluj-Napoca, Romania, 7-9 April 2022 •

SUPPLEMENT



Societatea Română de Obstetrică și Ginecologie



Societatea Română de Ultrasonografie în Obstetrică și Ginecologie



Societatea Română de HPV



Societatea de Uroginecologie din România



Societatea de Endometrioză și Infertilitate Est-Europeană



JOURNAL FOR CONTINUING MEDICAL EDUCATION

Year X • No. 35 (1) March 2022, Suppl. 1 • DOI: 10.26416/Gine.35.1.2022

ISSN 2457-5666; ISSN-L 2457-5666

Ginecologia

ro



FOTO: SHUTTERSTOCK

EDITOR-IN-CHIEF
Professor Radu VLĂDĂREANU

DEPUTY EDITOR-IN-CHIEF
Professor Daniel MUREȘAN
Lecturer Mihai MITRAN

EDITORIAL BOARD

Professor Victoria ARAMĂ; Dr. Diana BADIU;
Assoc. Professor Costin BERCEANU;
Assoc. Professor Roxana BOHÎLȚEA; Professor Elvira BRĂȚILĂ;
Professor Monica CÎRȘTOIU; Assis. Professor Ciprian COROLEUCĂ;
Assoc. Professor Camelia DIACONU;
Assoc. Professor Alexandru FILIPESCU;
Professor Claudia MEHEDINTU; Professor Marius MOGA;
Lecturer Claudiu MOLNAR; Assoc. Professor Dan NĂVOLAN;
Assoc. Professor Anca Maria PANAITESCU; Professor Liana PLEȘ;
Assoc. Professor Luminița PADURARU; Professor Cătălina POIANĂ;
Dr. Simona POPEȘCU; Lecturer Anca Angela SIMIONESCU;
Assoc. Professor Ana M.A. STĂNESCU;
Professor Simona VLĂDĂREANU

MIDWIVES AND NURSES SECTION COORDINATOR
Licensed Midwife Daniela STAN

SCIENTIFIC COMMITTEE

Nurse responsible for general care Carmen MAZILU
(President of OAMGMAMR, Bucharest subsidiary)
Nurse responsible for general care Anghelușa LUPU
(Vice-president of OAMGMAMR, Bucharest subsidiary)
Licensed assistant chief Anca MAREȘ
(Vice-president of OAMGMAMR, Bucharest subsidiary)
Nurse responsible for general care Corina GAGIU
(OAMGMAMR secretary, Bucharest subsidiary)

INTERNATIONAL EDITORIAL COMMITTEE

Professor Aris ANTSAKLIS (Athens, Greece)
Professor Frank CHERVENAK (NY, USA)
Professor Vincenzo D'ADDARIO (Bari, Italy)
Professor Valentin FRIPTU (Chișiinău, Republic of Moldova)
Professor Eberhard MERZ (Frankfurt/Main, Germany)
Professor Gianni MONNI (Cagliari, Italy)
Professor Milan STANOJEVIĆ (Zagreb, Croatia)
Professor Liliana VOTO (Buenos Aires, Argentina)
Professor Ivica ZALUD (Honolulu, USA)

DTP

Ioana BACALU

PROOFREADING

Florentin CRISTIAN

PHOTO PROCESSING

Radu LEONTE



CEO

Simona MELNIC

DEPUTY CEO MULTICHANNEL & EVENTS MANAGER

Lavinia SIMION

EDITORIAL MANAGER

Oana NEACȘU

SALES MANAGER

Mircea TOMESCU

ADMINISTRATIVE MANAGER

Dana STUPARIU

SUBSCRIPTIONS MANAGER

Cristina GRECU, Amelia SAVU
abonamente@medichub.ro

COMPANY ADDRESS:

MEDICHUB MEDIA SRL
Green Gate Office Building, 22 Tudor Vladimirescu Blvd.,
11th Floor, District 5, 050883, Bucharest, Romania
Phone: (031) 425.40.40, Fax: (031) 425.40.41
E-mail: redactia@medichub.ro
www.medichub.ro

Copyright © 2022 MEDICHUB MEDIA, S.R.L.
Drepturile de autor pentru articolele și fotografiile
publicate aparțin exclusiv MEDICHUB MEDIA, S.R.L.
Reproducerea, totală sau parțială, și sub orice formă,
tipărită sau electronică, sau distribuția materialelor
publicate se face numai cu acordul scris al Editurii.

ISSN 2344-2301

e-ISSN 2457-3566

ISSN-L 2344-2301

Responsabilitatea asupra conținutului original al materialelor
aparține în întregime autorilor. Persoanele intervievate
răspund de conținutul declarațiilor lor, iar utilizatorii
spațiului publicitar, de informațiile incluse în machete.

SUMMARIES OF THE 12TH NATIONAL CONFERENCE
OF THE ROMANIAN SOCIETY OF OBSTETRICS AND GYNECOLOGY,
CLUJ-NAPOCA, ROMANIA, 7-9 APRIL 2022

**Oligoamnios following SARS-CoV-2 vaccine or infection-induced? Case report
and literature review 8**

Cătălina Alexandrescu, Cosmina Catoiu, Romina-Marina Sima, Liana Pleș

**Optimization-based treatment guidelines in women with pelvic organ
prolapse 8**

Diana Badiu, Roxana Penciu, Vlad I. Tica

Transplacental transmission of hepatitis C virus infection 9

Elena-Iuliana-Anamaria Berbecaru, Anca-Maria Istrate-Ofițeru, Rodica Daniela Nagy,

Cătălina Rănescu, Alina-Iuliana Enache, Andreea Vochin, Ștefan-Gabriel Ciobanu,

Ionuț-Daniel Băluță, Laurențiu Mihai Diră, Ileana Drocaș, George-Lucian Zorilă,

Dominic-Gabriel Iliescu

Assessment, diagnosis and management in endometrial proliferations 10

Costin Berceanu, Flavia Andreea Dincă, Marinela Ionela Raicea, Oana Grigorie, Diana Mitran

**Management and outcome of preterm prelabor rupture
of membrane (PPROM) 10**

Roxana Bohîlțea, Ana-Maria Cioca, Ioniță Ducu, Bianca Mihai

**Project of national implementation of screening for cervical and breast
cancer 11**

Roxana Elena Bohîlțea

Fetal anomalies revealed late in pregnancy 11

Dan Boitor-Borza, Gheorghe Cruciat, Georgiana Nemeti, Adelina Staicu, Alexandra Centea,

Daniel Mureșan

**The impact of social support on the mental health of patients who gave
birth during the COVID-19 pandemic 12**

Dan Borza Boitor, Daiana Ștețco, Roxana Constantin, Daniel Mureșan

**Long-term results in radical vaginal trachelectomy with laparoscopic
lymphadenectomy for cervical cancer 12**

Petre Cornel Brătîlă, Elena Nadă, Anca Manta

**Vaginal reconstruction with acellular biologic mesh in congenital vaginal
malformations 13**

Petre Cornel Brătîlă, Elena Nadă, Anca Manta

**The distance between the embryo and the yolk sac in correlation
with the serum level of placental growth factor: how reliable is it? 13**

Carmen Elena Bucuri, Răzvan Ciortea, Valentin Oprea, Doru Diculescu, Andrei Mihai Măluțan,

Renata Nicula, Adelina Clim, Viorela Elena Suci, Maria Patricia Rada, Dan Mihu

Heart tumors and rhabdomyomas 14

Alina Mihaela Călin, Cătălin Herghelegiu

Epidemiological aspects of SARS-CoV-2 infection in the Obstetrics-Gynecology Clinic of Sibiu	14
Radu Chicea	
The challenges of diagnosis and management of pregnancy associated with neurofibromatosis and Becker nevus syndrome. Report of a unique clinical case	15
Monica Mihaela Cîrstoiu, Natalia Țurcan, Anca Simionescu, Octavian Munteanu, Delia Grădinaru-Fometescu, Florina Nedelea	
Therapeutic management in advanced vulvar cancer	15
Bogdan Cătălin Coroleucă, Ciprian-Andrei Coroleucă, Andrei Manu, Alexandra Baușic, Diana-Elena Comandașu, Diana Mihai, Elvira Brătîlă	
Laparoscopic hysterectomy in Bega clinic from Timișoar	16
Marius Craina, Elena Bernad, Carmen Silaghi, Adelina Erimescu, Zoran Popa, Radu Neamțu, George Dahma	
The value of laparoscopic surgery in adnexal torsion	16
Marius Craina, Elena Bernad, Carmen Silaghi, Adelina Erimescu, Zoran Popa, Radu Neamțu, George Dahma	
Preeclampsia – screening, monitoring and clinical cases	17
Oana Eliza Crețu, Alina-Alexandra Dîrlău, Cristian Poalelungi, Dragoș Dobrițoiu, Iuliana Ceaușu	
Implications of hypertriglyceridemia-triggered acute pancreatitis during pregnancy	17
Gheorghe Cruciat, Georgiana Nemeti, Iulian Goidescu, Andreea Florian	
Comparison of demographic and ultrasound regression formulas for hypertensive pregnancy complications	18
Voicu Dașcău, Gheorghe Furău, Luminița Pilat, Cristina Onel, Adeline Horga, Iman Ait El Haj, Cristina Popescu, Maria Pușchița	
Statistical comparison of regression formulas for hypertensive pregnancy complications	18
V. Dașcău, Gheorghe Furău, Luminița Pilat, Cristina Onel, Adeline Horga, Iman Ait El Haj, Cristina Popescu, Maria Puschița	
The benefits of HPV vaccination beyond prophylaxis	19
D. Diculescu, D. Mihu, C. Iuhaș, Renata Nicula, Carmen Bucuri, A. Măluțan, R. Ciortea, Marina Dudea, Maria Rada, Daria Pop	
Thrombophilia-related and placenta-mediated pregnancy complications	19
Flavia Andreea Dincă, Marinela Ionela Raicea, Elena Loredana Ciurea, Oana Grigorie, Diana Mitran, Costin Berceanu	
COVID-19 pandemic effects on maternal follow-up and anemia as a pregnancy complication – experience from a tertiary care center	20
Alina-Alexandra Dîrlău, Oana-Eliza Crețu, Delia-Cristina Angelescu, Alice Ionescu, Cristina-Ioana Iancu, Djamilia Djafer, Cristian Poalelungi, Dragoș Dobrițoiu, Iuliana Ceaușu	
Why use an ultrasound evaluation in the labor ward?	20
Roxana-Cristina Drăgușin, Laurențiu Dîră, Maria Șorop-Florea, Marina Dinu, Lucian Zorilă, Rodica Nagy, Alina Enache, Anca-Maria Istrate-Ofițeru, Dominic-Gabriel Iliescu	
Vulvar cancer – a case report of a patient who presented vulvar squamous cell carcinoma with rapid disease progression	21
Mihai Cristian Dumitrașcu, Florica Șandru, Olga Scaletchi, Răzvan Fodoriu	
The importance of the three-vessel trachea view in prenatal screening	21
Roxana Gireadă, Răzvan Socolov, Elena Mihălceanu, Luiza Guzgă	
Advanced breast carcinoma diagnosed during pregnancy	22
Iulian Goidescu, Georgiana Nemeti, Andreia Preda, Adelina Staicu, Gabriela Caracostea, Mihai Surcel, Gheorghe Cruciat, Daniel Mureșan	
Immunomodulation in genital HPV infections – in between mirage and real benefit	22
Corina Grigoriu, Gina Ionescu, Andra Bălan, Irina Horhoianu	

Large loop excision of the transformation zone: “see and treat” or “think twice about it”?	23
Corina Grigoriu, Gina Ionescu, Andra Bălan, Irina Horhoianu	
Role of pulse oximetry in the screening of congenital heart diseases	23
Monica Gabriela Hășmășanu, Liliana Kalisch, Margareta Pârvănoiu, Melinda Matyas, Ligia Blaga, Gabriela Zaharie	
The late preterm infants – a continuous challenge	24
Monica Gabriela Hășmășanu, Bianca Marcela Voina, Melinda Matyas, Ligia Blaga, Dan Boitor, Andrada Mădălina Petyan, Gabriela Zaharie	
Inherited thrombophilia in pregnancy: a cause of postpartum hemorrhage?	24
Ciprian Ilea, Demetra Socolov, Irina Stoian	
The importance of diagnosis of uterine congenital malformations in pregnant patients	25
Alice Ionescu, Delia Angelescu, Alina-Alexandra Dîrlău, Oana-Eliza Crețu, Cristina Iancu, Alexandru Ciulcu, Iuliana Ceaușu	
Obstetrical soft tissue trauma during vaginal delivery in the Romanian adolescent population – a multicentric comparative study with adult population	25
Crîngu Ionescu, Alexandra Matei, Mihai Dimitriu, George Roșu, Dan Năvolan, Cristian Furău	
Update on maternal immunization excluding COVID-19 vaccine	26
Costin Alexandru Ispas, Antoine Edu, Elena Ispas, Dimitrie Nanu	
Induction of labor using the Foley balloon in patients with post-caesarean scar uterus	26
Victor Juncu, Camille Olivier, Călin Gheorghe Oancea, Mihai Emil Căpîlna	
Considerations on COVID-19 pregnancy: a case series during 2020-2021 in Prahova region, southern Romania	27
Ioan Bogdan Luchian, Ali Karaouni, Călin Boeru, Letiția Gorăneanu	
Preeclampsia – a myriad of miRNAs	27
Andrei Mihai Măluțan, Doru Diculescu, Răzvan Ciortea, Cristian Iuhaș, Radu Florin Mocan-Hognogi, Carmen Bucuri, Maria Rada, Marina Ducea, Viorela Suci, Miha Dan	
The role of adipokines in uterine contractility	28
Dan Miha, Doru C. Diculescu, Răzvan Ciortea, Andrei M. Măluțan, Ciprian G. Porumb, Mihaela D. Oancea, Renata L. Nicula, Cristina M. Prodan	
Maternal HIV infection – prematurity risk factor?	28
Mihai Mitran, Octavia Velicu, Elvira Brătîlă	
Magnesium sulphate as a neuroprotective agent in imminent risk for preterm delivery – a challenge for obstetrical practice	29
Marius Alexandru Moga, Sabina Ioana Bădilă, Oana Dimienescu, Cezar Podașcă, Cristian Arvătescu, Nicușor Bigiu, Andreea Bălan	
Pregnancy and neuromyelitis optica spectrum disorder (NMOSD) – a rare but challenging interaction	29
Georgiana Nemeti, Gheorghe Cruciat, Ioana Rotar, Gabriela Caracostea, Iulian Goidescu, Andreia Preda, Mihai Surcel, Dan Boitor, Daniel Mureșan	
Increase of the rates of premature birth and of caesarean section generated by COVID-19 in a tertiary maternity	30
Octavian Gabriel Olaru, Anca Daniela Stănescu, Ileana Conea, Denisa Oana Bălălau, Romina Marina Sima, Liana Pleș	
Cervical cancer screening in Romania: it is time to change	30
Gheorghe Peltecu, Anca Maria Panaitescu, Nicolae Gică	
Sentinel lymph node biopsy in malignant pathology of vulva	31
Gheorghe Peltecu, Anca Maria Panaitescu, Nicolae Gică	

Does birth improve the outcome of pregnancies with mild or severe SARS-CoV-2 infection?	
Our experience	31
Liana Pleș, Romina-Marina Sima, Tina-Ioana Bobei, Georgiana Răducanu, Anca-Daniela Stănescu	
The impact of SARS-CoV-2 pandemic on obstetrics-gynecology residents' education	32
Liana Pleș, Romina-Marina Sima, Ioana Ciupitu, Anca-Daniela Stănescu	
The hysteroscopic management of abnormal uterine bleeding	32
Cristian Poalelungi, Iuliana Ceaușu, Nicoleta Alina Mareș, Maria Videnie	
Adolescent pregnancy: complications and birth rates	33
Daria Maria Pop, Renata Nicula, Cezarin Todea, Ciprian Porumb, Hognogi Radu Mocan, Ligia Blaga, Dan Mihiu, Doru Diculescu	
The usefulness of the SPIKES protocol in obstetrician-couple communication in case of detecting intrauterine fetal malformations	33
Dragoș Popescu, Radu Chicea, Iulia Lazăr	
Postoperative period management in patients after abdominal birth	33
Oleg Potacevski	
The influence of locoregional anesthesia on the evolution of the postoperative period	34
Oleg Potacevski	
Fetal shunt placement	34
Hadi Rahimian, Carmen Cristea, Oana Zaharia, Camelia Miron	
Twin-twin transfusion syndrome management	35
Hadi Rahimian, Carmen Cristea, Oana Zaharia, Camelia Miron	
Indoor environmental quality assessment in a historical building – a maternity field-study in North-West Romania	35
Tania Rus, Gheorghe Cruciat, Georgiana Nemeti, Iulian Goidescu, Florin Vasile Domnița, Daniel Mureșan	
Cervical remodeling in preterm birth. Timing, diagnosis and actual therapeutic opportunities	36
Manuela Russu, Adriana Ghelme, Ruxandra Stănculescu, Șerban Nastasia	
High-risk HPV involvement in post-therapeutic CIN2+/CIN3+ recurrence	36
Manuela Russu, Cătălina Savu, Ruxandra Stănculescu, Anca Popa, Șerban Nastasia	
The place of colposcopy in modern gynecology. Indications, logistics and procedures	37
Ioan Sas, Larisa Tomescu, Maia Todorov	
Second victims in obstetrics – a literature review	37
Ioan Sas, Nicoleta Nicolae, Denis Șerban, Larisa Tomescu	
Current clinical and therapeutic challenges in HELLP syndrome	38
Romina-Marina Sima, Gabriel Gorecki, Andreea Marcu, Liana Pleș	
The pathological characteristics of myometrium in uterine atonia – case report	38
Romina-Marina Sima, Cristina-Diana Pomană, Liana Pleș, Cătălina Alexandrescu	
Management challenges of psoriasis in pregnancy	39
Anca Angela Simionescu	
Epidemiology of clavicle fracture at birth in Romania.....	39
Anca Angela Simionescu, Vlad Dima	

Placental pathologic changes in gestational diabetes mellitus and implications for fetal outcome	40
Adelina Staicu, Héloïse Pelczar, Camelia Albu, Ioana Cristina Rotar, Andreia Preda, Roxana Constantin, Iulian Goidescu, Dan Boitor-Borza, Mihai Surcel, Gheorghe Cruciat, Daniel Mureșan	
Update of the epidemiology and ultrasound diagnosis criteria able to predict before delivery	
placenta accreta spectrum disorders	41
Ruxandra Stănculescu, Manuela Russu, Elvira Brătîlă, Vasilica Bausic, Oana Toader, Ciprian Coroleucă, Alexandra Bausic, Patru Papuc, Ramona Dragomir	
The impact of SARS-CoV-2 infection on fluid-coagulant status of pregnant women	41
Anca-Daniela Stănescu, Romina-Marina Sima, Liana Pleș	
Short-term disease recurrence after loop electrosurgical excision procedure (LEEP) – retrospective study	42
Irina Stoian, Ciprian Ilea, Demetra Socolov	
Magnesium sulphate and fetal neuroprotection – when, how, why and why not?	42
Viorela-Elena Suci, Răzvan Ciorte, Carmen Elena Bucuri, Andrei Măluțan, Radu Mocan-Hognogi, Maria Rada, Marina Ducea, Adelina Clim, Dan Mih	
Therapeutic solutions for infertile “poor responder” patients	43
Mihai Surcel	
Ultrasound aspects of uterine scars after caesarean section in patients with retroverted uterus versus patients with anteverted uterus	43
Lucian Șerbănescu, Paris Ionescu, Diana Badiu, Dan Costea	
Incidence of premature birth threat in patients with fibromatous uterus	43
Lucian Șerbănescu, Paris Ionescu, Diana Badiu, Dan Costea	
The impact of autoimmune, inflammatory and metabolic disorders on placental inflammation and perinatal outcomes	44
Atakan Tanacan	
Conotruncal congenital heart diseases: prenatal diagnosis and management	44
Atakan Tanacan	
Management of gastroesophageal reflux during pregnancy and lactation	45
Vlad Tica, Roxana Penciu, Liliana Steriu, Silvia Izvoranu, Corina Nour, Mihai Banariu, Diana Mocanu, Diana Badiu, Irina Tica	
Vaginal senescence in menopause	45
Vlad Tica, Silvia Izvoranu, Mihai Banariu, Roxana Penciu, Liliana Steriu, Corina Nour, Diana Mocanu, Diana Badiu, Irina Tica	
Is there an endometrium-ovarian cancer link?	46
Vlad Tica, Liliana Steriu, Diana Badiu, Roxana Penciu	
Correlation between genital prolapse and obesity	46
Mihaela Camelia Tirnovanu, Roxana Mihaela Corduneanu, Andreea Crudu, Vlad Gabriel Tirnovanu, Bogdan Toma, Alexandra Iov	
Clinical and prognostic implications regarding the molecular classification of invasive breast carcinoma	47
Oana Toader, Alexandra Olaru, Mădălina Mârza, Ramona Dragomir, Tiberiu Georgescu	
Current concepts and etiopathogenic considerations in the diagnosis of pelvic serous tumors	47
Oana Toader, Ramona Dragomir, Mădălina Mârza, Ramona Mircea-Vicol, Tiberiu Georgescu	
Multigene NGS panels and cancer susceptibility syndromes – a study report	48
Radu-Ioan Ursu, Diana Prepeliță, Georgiana Ilie, Cristina Dragomir, Iuliana Chelu, Grațiela Chelu, Simona Filimon, Mihaela Popa, Bianca Basangiu, Andreea Ionescu, Andra Perioc, Ioana Patricia Riurean, Antoanela Curici	

Particularities of acute pyelonephritis in pregnant women	48
Cristian Vasilachi	
Treatment of chronic endometritis in cases with implantation failure after <i>in vitro</i> fertilization procedure	49
Cristian Văduva, Laurențiu Dîră, Sidonia Săndulescu, Alice Văduva	
Progesterone-follicle ratio is better correlated than blood progesterone level with <i>in vitro</i> fertilization results	49
Cristian Văduva, Sidonia Săndulescu, Laurențiu Dîră, Alice Văduva	
The outcome of structural heart defects diagnosed in the first trimester of pregnancy	50
Alina Veduță	
Management of unexpected <i>placenta accreta</i> spectrum after vaginal delivery – case report	50
Octavia Velicu, Cristina Popescu, Eugen Tănase	
The new Enzian classification and the correlation between preoperative evaluation of deep infiltrating endometriosis and intraoperative findings	51
Simedrea Voicu	
Methods of evaluating the colorectal and urinary surgery quality in colorectal and urinary surgery regarding advanced deep infiltrating endometriosis	51
Simedrea Voicu	
Low-impact laparoscopy, ERAS and multimodal analgesia in endometriosis advanced surgery	52
Simedrea Voicu	

Oligoamnios following SARS-CoV-2 vaccine or infection-induced? Case report and literature review

Cătălina Alexandrescu¹, Cosmina Catoiu², Romina-Marina Sima^{2,3}, Liana Pleș^{2,3}

1. Medlife Memorial Hospital, Bucharest

2. "Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

3. Department of Obstetrics and Gynecology, "Sf. Ioan" Clinical Hospital, Bucur Maternity, Bucharest, Romania

Introduction. The literature reports and recommendations are clear about the SARS-CoV-2 vaccine safety during pregnancy. The majority of pregnant women have a positive attitude towards SARS-CoV-2 vaccine, but some of them are still sceptic due to potential side effects. **Materials and method.** We searched the literature about the fetal consequences of SARS-CoV-2 vaccine following its usage during pregnancy. The keywords were: "SARS-CoV-2 vaccine", "pregnancy", "side effects", and "oligoamnios". The research was initiated by a unique case that was investigated in our clinic and we present it. **Results.** The literature reports a reduce number of studies regarding the SARS-CoV-2 vaccine impact during pregnancy. Our research revealed 78 publications. It is already established that pregnant individuals are more likely to develop severe COVID-19 and a significant number of pregnancy complications have been observed in COVID-19 patients. The majority of studies explore the role of the maternal antibody response to SARS-CoV-2

during pregnancy and the transfer of maternal antibodies from COVID-19 infected mothers to the fetus. We report a case with severe oligoamnios developed after SARS-CoV-2 vaccine in the third trimester of pregnancy. The placental insufficiency and uncertain fetal status imposed caesarean section. The maternal and fetal outcome were favorable. Antibody serological investigation revealed for both mother and new born high titers of nucleocapsidar and antispikes antibodies, significant for suspecting that the mother had the infection during pregnancy, and she also was vaccinated. The pathological exam of the placenta revealed important but nonspecific inflammatory response. **Conclusions.** The COVID-19 pandemic is still affecting the general population, including pregnant women. There are isolated cases of SARS-CoV-2 vaccine side-effects in pregnancy. Even so, women should be encouraged to receive the vaccine during pregnancy in order to avoid the severe consequences of COVID 19.

Keywords: oligoamnios, vaccine, SARS-COV-2

Optimization-based treatment guidelines in women with pelvic organ prolapse

Diana Badiu, Roxana Penciu, Vlad I. Tica

Department of Obstetrics and Gynecology, Institute of Doctoral Studies, Doctoral School of Medicine, Faculty of Medicine, "Ovidius" University of Constanța; "Sf. Apostol Andrei" Emergency Clinical County Hospital, Constanța, Romania

Pelvic organ prolapse (POP) represents a common gynecological disorder which affects the women's quality of life (QoL). Nowadays, many treatment options for women with POP have been developed in clinical practice. When discussing treatment options with POP patients, gynecologists make recommendations based more on clinical experience than on specific clinical guidelines. Our objective in the present review is to

identify optimization-based treatment guidelines for women with POP to serve as a reference in daily practice. We address the question related to the optimization methods for specific treatment guidance, which maximize the results and the patient's expectation regarding their future QoL.

Keywords: pelvic organ prolapsed, treatment optimization, guidelines, patient's expectation

Transplacental transmission of hepatitis C virus infection

Elena-Iuliana-Anamaria Berbecaru³, Anca-Maria Istrate-Ofițeru^{1,2,3}, Rodica Daniela Nagy³, Cătălina Rănescu³, Alina-Iuliana Enache³, Andreea Vochin³, Ștefan-Gabriel Ciobanu³, Ionuț-Daniel Băluță³, Laurențiu Mihai Diră³, Ileana Drocaș^{2,3}, George-Lucian Zorilă^{2,3}, Dominic-Gabriel Iliescu^{2,3}

1. Department of Histology, University of Medicine and Pharmacy of Craiova, Romania

2. Department of Obstetrics and Gynecology, University of Medicine and Pharmacy of Craiova, Romania

3. Second Department of Obstetrics and Gynecology, Emergency County Hospital of Craiova, Romania

Introduction. Hepatitis C virus (HCV) can cause acute or chronic hepatitis. HCV can be spread through inadequate sterilization or reuse of syringes and needles, blood transfusion, injecting drug use, sexual intercourse, and vertical transmission from the mother to the fetus, although the last two are very rare. Vertical transmission of HCV occurs in rare cases (5-8%), more often during birth, and seldom transplacental, before birth. Women infected with HCV are more predisposed to give birth to preterm babies or low birth weight babies, or babies with congenital anomalies. **Materials and method.** We present the case of a 42-year-old woman, gravida 2, para 2, with a 32-33-week pregnancy, with not investigated deaf-mutism, who came in the emergency room for preterm labor contractions. At the ultrasound exam, we noticed fetal ascites, polyhydramnios and echogenic fetal bowel. The patient was admitted to our clinic for further investigations. We recommended an amniocentesis and a genetic consultation. The mother was tested for the TORCH panel, including HCV. The anti-HCV antibodies were present. In our service, the mother was also diagnosed with gestational diabetes. Following the clinical and paraclinical investigations performed, we decided to perform fetal cordocentesis, paracentesis and amniocentesis. Under ultrasound guidance, we performed cordocentesis and obtained approximately 8 ml of fetal blood and sent it for the following biological tests: albumin, direct bilirubin, indirect bilirubin, total bilirubin, blood count, blood glucose, direct Coombs test, HCV RNA, IgM antibodies for cytomegalovirus/toxoplasmosis/rubella/herpes simplex type 1 and 2. For the second puncture, the catheter was introduced at the level of the fetal peritoneal cavities under ultrasound guidance. We extracted 160 ml of fetal ascites fluid and sent it for cytological and biochemical analysis (leukocytes, red blood cells, glucose, albumin, creatinine, urea, proteins, Rivalta test, cytological examination). We also performed amniocentesis and collected 40 ml of clear

amniotic fluid. We used it to detect an infection with herpes simplex virus, Epstein-Barr virus, or parvovirus. **Results.** The biochemical tests and blood count performed from the fetal blood were within normal range. The direct Coombs test was negative. HCV RNA was detected (251 UI/mL). IgM antibodies for cytomegalovirus/toxoplasmosis/rubella/herpes simplex type 1 and 2 were all absent, and antibodies for Epstein-Barr virus and parvovirus B19 were also absent. The cytological exam of the ascites fluid showed smears with hypercellularity: very frequently lymphocytes, rare isolated mesothelial cells within normal cytological limits, rare polymorphonuclear leukocytes, and rare red blood cells. The biochemical tests performed from the fetal ascites fluid showed: red blood cells (3000/mm³), leukocytes 835/mm³, glucose 57 mg/dL, albumin 1.90 g/dL, creatinine 0.53 mg/dL, urea 19 mg/dL, proteins 3.4 g/dL, positive Rivalta test. HCV RNA was detected. IgM antibodies for cytomegalovirus/toxoplasmosis/rubella/herpes simplex type 1 and 2 were all absent. After fetal diagnosis, our colleagues from the infectious disease department recommended parenteral/oral hepatoprotective treatment for the mother; there was no possibility of antiviral treatment of the fetus. After performing the paracentesis of the fetus, the amniotic fluid was in the normal range, the fetal ascites were in small quantity with ultrasonographic hepatosplenomegaly aspect and normal echogenicity of the bowel. Almost seven weeks after the paracentesis, the patient gave birth to a baby, gender male, weight 2880 g, with an Apgar score of 7, which was handed over to the neonatology service, for further investigations. **Conclusions.** In rare cases, HCV can be transmitted vertically from the mother to the unborn baby. Cordocentesis and paracentesis helped us diagnose the fetus before birth and improve his prognosis after birth.

Keywords: ascites, HCV infection, cordocentesis, paracentesis, amniocentesis

Assessment, diagnosis and management in endometrial proliferations

Costin Berceanu¹, Flavia Andreea Dincă², Marinela Ionela Raicea², Oana Grigorie³, Diana Mitran³

1. Department of Obstetrics and Gynecology, County Emergency Clinical Hospital Craiova; University of Medicine and Pharmacy of Craiova, Romania

2. Department of Obstetrics and Gynecology, County Emergency Clinical Hospital Craiova, Romania

3. Doctoral School of Medicine, University of Medicine and Pharmacy, Craiova, Romania

Endometrial hyperplasia (EH) represents an abnormal endometrial proliferation, characterized by an increase in the gland/stroma ratio >50%. EH is the most common cause of vaginal bleeding, both before and after menopause. Two systems exist for classifying EH: the World Health Organization (WHO) system and the endometrial intraepithelial neoplasia (EIN) system. According to EIN criteria, endometrial proliferative lesions include benign, reactive EH, intraepithelial endometrial neoplasia and endometrial carcinoma. Ultrasonography (US) is the main imaging technique for assessing the pelvis, both in terms of diagnosis and patient compliance. Two-dimensional US, through the sagittal and transverse planes, offers some limitations in the examination of endometrial pathology, while 3D examination adds the coronal plane, allowing the evaluation of the entire volume and continuity of angular structures. Sonohysterography may also be included among endometrial imaging options. Diagnostic and operative hysteroscopy, through the direct endometrial approach, represents the suc-

cessive and complementary stage in the diagnosis and therapy of endometrial proliferations. The succession of these two explorations, together with the experience of the examiner, can approach the accuracy of the pathological examination, with a concordance of 78-100% depending on the technique and the type of lesion. The correlation of US examination with hysteroscopy data and later with the surgical pathology diagnosis outlines the complete picture of the exploration of benign or malignant endometrial proliferations, both in terms of study and evaluation of the endometrium as an implantation marker, in terms of fertility status and subsequent proliferative potential in peri- and post-menopause. This paper aims at an integrated presentation of the endometrial proliferations through the perspective of imaging techniques and those of direct endometrial approach in correlation with the surgical pathology diagnosis and therapeutic management of these lesions.

Keywords: endometrium, hyperplasia, ultrasound, hysteroscopy, pathology

Management and outcome of preterm prelabor rupture of membrane (PPROM)

Roxana Bohilțea^{1,2}, Ana-Maria Cioca¹, Ioniță Ducu², Bianca Mihai²

1. Department of Obstetrics and Gynecology, "Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

2. Department of Obstetrics and Gynecology, University Emergency Hospital, Bucharest, Romania

The management of prelabor rupture of membrane (PPROM) is one of the most discussed issues in perinatal medicine. The guidelines recommend to manage the pregnancy expectantly whenever this is possible, taking into account certain factors, such as gestational age, presence or absence of maternal/fetal infection, presence or absence of labor, fetal presentation, fetal well-being, and cervical status. Our objective is to demonstrate the benefits of expectantly management. We undertook a retrospective observational analytical study in which we included PPRM pregnancy between 2015 and 2020. We followed the evolution of premature newborn using several parameters (neonatal death, respiratory distress syndrome, hypoxia, intraventricular hemorrhage, hypo-

glycemia, necrotizing enterocolitis, neonatal sepsis), originated from both PPRM pregnancies that were managed expectantly and PPRM that were delivered short time after admission. In PPRM pregnancies managed expectantly we included the patients to whom there had been administered tocolysis, corticosteroids and antibiotics, so the delivery was postponed. We observed a significant decrease in neonatal complication in pregnancies that were prolonged. We concluded that the expectantly management of pregnancies with preterm prelabor rupture of membrane leads to a statistical benefit, decreasing the incidence of fetal and maternal complications.

Keywords: PPRM, neonatal, pregnancy, premature newborn, expectantly

Project of national implementation of screening for cervical and breast cancer

Roxana Elena Bohilțea

Department of Obstetrics and Gynecology, "Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

The development of the Romanian society and medical education in the last 30 years will allow the medical system to move from the area of medical care of advanced cases of the disease to the area of primary and secondary prevention, in accordance with global efforts, increasing the duration and quality of life of individuals in populations whose health will be so greatly improved. The current screening of breast and cervical cancer in Romania remains predominantly opportunistic, inconsistent and uncoordinated. A personal project based on the analysis of the implementation problems of national screening programs in Romania supports the need to create specialized regional screening centers with financial and operational autonomy, with dedicated staff, equipment and standardized procedures, capable to ensure the development of population screening programs, according to the requirements and quality standards of the European Union. The national program should be remodeled, focusing on the rural population and, in particular, the western, southern and south-western regions, which still

maintains major mortality differences from the national level. Another essential prerequisite for the national program is to ensure the population character of screening, which is in line with European guidelines and domestic legislation guaranteeing universal access to preventive services for all citizens. Individual invitations *via* screening register or database must be linked to the population record database, cancer register, rare diseases register and mortality database. A fixed appointment date must be included in the invitation and a service continuum for positive cases has to be designed. I propose a radical change in the national approach to screening, through the full financial-operational dichotomization of medical services for therapy and screening. The establishment of these autonomous centers will open the way and provide the material basis for other screening programs, like the one for colorectal cancer that is already a requirement of the European Union, for lung and prostatic cancer screening programs that are currently evaluated.

Keywords: screening, cervical cancer, breast cancer

Fetal anomalies revealed late in pregnancy

Dan Boitor-Borza, Gheorghe Cruciati, Georgiana Nemeti, Adelina Staicu, Alexandra Centea, Daniel Mureșan

First Clinic of Obstetrics and Gynecology, County Emergency Clinical Hospital Cluj-Napoca; "Iuliu Hațieganu" University of Medicine and Pharmacy, Cluj-Napoca, Romania

The diagnosis of fetal anomalies in the first trimester of pregnancy could be a challenging task; mostly, they are seen on the occasion of the second-trimester ultrasound. However, some fetal anomalies reveal only late in pregnancy. Finding an anomaly in the third trimester is an unpleasant surprise both for the parents and sonographer, as it can generate ethical, medical and even legal problems. We present some of the fetal anomalies that can be diagnosed by ultrasound only in the late mid-trimester or in the third trimester of pregnancy. Cerebral tumors, ventriculomegaly, cardiac rhabdomyoma, aneurysm of the left ventricle, renal anomalies and digestive anomalies correlated with hydramnios are discussed. We insist on the necessity

of correlation with magnetic resonance imaging findings in difficult cases; thus, accurate diagnosis, better management and precise prognosis could be provided. **Conclusions.** Sonographers must be aware that certain fetal anomalies could be seen only late in pregnancy, as present earlier but impossible to be observed at that time, or developed in the second half of pregnancy. On the other hand, parents must be informed that some pathologies develop in the first and second trimesters, and can be diagnosed by ultrasound only in the third trimester. Neonatologists and other staff that provide medical care to the affected babies have to know that certain anomalies develop late in pregnancy.

Keywords: fetal anomalies, ultrasound, late diagnosis

The impact of social support on the mental health of patients who gave birth during the COVID-19 pandemic

Dan Borza Boitor, Daiana Ștețco, Roxana Constantin, Daniel Mureșan

First Clinic of Obstetrics and Gynecology, County Emergency Clinical Hospital Cluj-Napoca; "Iuliu Hațieganu" University of Medicine and Pharmacy, Cluj-Napoca, Romania

Introduction. The postnatal period is for a woman a time of great vulnerability to become mentally unwell, and postpartum mood disorders are likely exacerbated by the COVID-19 pandemic. This retrospective study aims to understand the impact of social support on mental health of patients who gave birth during the COVID-19 pandemic. **Method.** The participants were 163 women (19-40 years old) recruited from hospital to complete an online survey between May and July 2021. We assessed depression, anxiety and perceived stress using validated questionnaires. Social support was assessed in relation to mental health. **Results.** Linear regression analysis

identified that social support has been a small predictor, explaining 10% of the variation ($\beta=-32$, $p<0.001$) in postnatal depression scores, 19% of the variation in anxiety scores ($\beta=-44$, $p<0.001$), and 5% of the variation in perceived stress scores ($\beta=-234$, $p<0.003$). **Conclusions.** This study shows that social support had a small impact on the mental health of the patients who gave birth during the COVID-19 pandemic. Contrary to expectation, social support was found not to be a significant predictor of postnatal depression, anxiety and perceived stress.

Keywords: COVID 19 pandemic, mental health, social support

Long-term results in radical vaginal trachelectomy with laparoscopic lymphadenectomy for cervical cancer

Petre Cornel Brătilă, Elena Nadă, Anca Manta

Gynecology Clinic, Euroclinic Minimally Invasive Surgery, "Regina Maria" Network, Bucharest, Romania

Introduction. Radical vaginal trachelectomy with laparoscopic pelvic lymphadenectomy is the proposed solution for patients diagnosed with incipient cervical cancer who desire to maintain their fertility, a procedure that can also be performed during pregnancy. **Materials and method.** A retrospective study was conducted in our clinic over a period of 15 years. We included 85 patients diagnosed with cervical cancer in early stages, who underwent conservative treatment, consisting of radical trachelectomy and radical pelvic lymphadenectomy or pelvic sentinel lymph node biopsy. Two cases (1.7%) in which conversion to open surgery was performed were excluded. Among the included patients, 17 (20.48%) had viable pregnancies. Two procedures (2.4%) were performed in patients during the first and second trimesters of pregnancy. The trachelectomy was performed vaginally in 97.59% of cases and laparoscopically in 2.4% of cases. The pelvic lymphadenectomy was performed laparoscopically in

100% of cases. All patients were staged from CIS to stage IB1 FIGO (tumor <2 cm). The average age was 31.93 years old. All patients had HPV-HR+. Local recurrence was recorded in only one case (0.83%). The postoperative follow-up was performed by clinical examination, cytology and imaging. **Discussion.** The conditions for performing the conservative treatment are described in the internal protocol. Radical vaginal trachelectomy offers many advantages compared to the abdominal approach: minimally invasive anatomical surgery, it preserves the vascularity of the remaining uterus, it avoids ureteral complications. **Conclusions.** Radical vaginal trachelectomy with pelvic lymphadenectomy or pelvic sentinel lymph node biopsy is a safe procedure suitable in the management of early cervical cancer in women who desire to preserve their reproductive function or in a pregnant patient.

Keywords: cervical cancer, fertility, conservative treatment

Vaginal reconstruction with acellular biologic mesh in congenital vaginal malformations

Petre Cornel Brătilă, Elena Nadă, Anca Manta

Gynecology Clinic, Euroclinic Minimally Invasive Surgery, "Regina Maria" Network, Bucharest, Romania

Introduction. Mayer-Rokitansky-Kuster-Hauser syndrome (MRKH) is a complex malformation of the female urogenital tract, including vaginal and uterine agenesis. Vaginal agenesis is also found in Morris syndrome. The aim of this paper is to present our experience in vaginal reconstruction. **Materials and method.** A retrospective study was conducted over a period of four years in our clinic. We included eight patients diagnosed with complex cervico-utero-vaginal malformations, in whom laparoscopically assisted vaginal neocolpoptosis with acellular biologic mesh was performed. Two patients (25%) had type II MRKH syndrome (associated with urinary tract malformations). Seven patients (87.5%) had 46XX karyotype, and patient (12.5%) had 46XY karyotype. The average age at diagnosis was 16.5 years old. The average age at the time of the intervention was 19.3 years old. The average operating time was 176 minutes and the average length of hospitalization was 72 hours. A long-term complication was registered, consisting in the stenosis of the vaginal graft. Seven pa-

tients (87.5%) had a normal sex life. The average length of the vagina was 9.4 cm. The postoperative follow-up was performed by clinical examination every 10 days, 30 days and every 3 months in the first year. **Discussion.** MRKH syndrome is an association of congenital defects of the Müllerian ducts. Depending on the association with other defects (urinary, intestinal etc.), it is divided into two types. The diagnosis is usually established at puberty, due to primary amenorrhea. There are several methods of vaginal reconstruction, using ileal, sigmoid, skin, muscle or acellular biologic mesh. The surgical outcome depends on the patient's postoperative cooperation. The main complication of this technique is vaginal stenosis. **Conclusions.** Vaginal reconstruction is a challenge in both gynecological and plastic surgery. Acellular biologic mesh neocolpoptosis is a viable method of treating vaginal agenesis, with good results regarding patient's satisfaction and sexuality.

Keywords: vaginal reconstruction, dermal matrix, malformations

The distance between the embryo and the yolk sac in correlation with the serum level of placental growth factor: how reliable is it?

Carmen Elena Bucuri^{1,2,3}, Răzvan Ciortea^{1,2}, Valentin Oprea^{1,3}, Doru Diculescu^{1,2}, Andrei Mihai Măluțan^{1,2}, Renata Nicula^{1,2}, Adelina Clim², Viorela Elena Suci², Maria Patricia Rada^{1,2}, Dan Miha^{1,2}

1. "Iuliu Hațieganu" University of Medicine and Pharmacy, Cluj-Napoca, Romania

2. "Dominic Stanca" Obstetrics-Gynecology Clinic, County Emergency Clinical Hospital, Cluj-Napoca, Romania

3. "Dr. Constantin Papilian" Military Emergency Clinical Hospital, Cluj-Napoca, Romania

Introduction. The etiology of embryonic demise is multifactorial, with chromosomal abnormalities being the most common (60%). The purpose of this study is to evaluate the correlation between a serum biomarker, placental growth factor (PIGF), and an ultrasonographic parameter, the distance between yolk sac (YS) and embryo (DYSE), in assessing the prognosis of pregnancy evolutivity in the first trimester. **Method.** The study is a case-control prospective analysis that includes two groups of patients: 81 patients with first-trimester pregnancy in evolution, and 89 patients with embryonic demise, all of the patients having amenorrhea between 6 and 11 weeks. The endovaginal ultrasonographic exploration was performed to evaluate the distance between the lower pole of the embryo and the yolk sac. From

each subject enrolled in the study, 20 ml of blood was collected for PIGF dosing. **Results.** Regarding the DYSE in the case group, lower values were observed compared to the control group, the difference being statistically significant. In the statistical analysis of serum PIGF values, statistically significant differences were observed between the two groups ($p < 0.0001$). **Conclusions.** DYSE has a high positive predictive value in identifying pregnancies with potentially reserved evolutivity, this study demonstrating that a $DYSE < 3$ mm may lead to an unfavorable evolution of pregnancy. The low serum level of PIGF is associated with an increased rate of nonviable embryos.

Keywords: embryonic demise, first trimester, ultrasonography

Heart tumors and rhabdomyomas

Alina Mihaela Călin¹, Cătălin Herghelegiu²

1. "Sf. Apostol Andrei" Emergency Clinical Hospital, Galați, Romania

2. "Polizu" Hospital, "Alessandrescu-Rusescu" National Institute for Mother and Child Health, Bucharest, Romania

Introduction/objective. Heart tumors are rare and represent 1-2% of all cardiac anomalies. The most common type of tumors encountered are rhabdomyomas (80-90% of cases), but there can also be teratoma, fibroma, myxoma, hamartoma, rhabdomyosarcoma and others. Fetal outcome depends on the tumor size, impairment of flow, and the occurrence of rhythm disturbances. However, rhabdomyomas tend to shrink after birth and even eventually disappear completely. **Method.** Rhabdomyomas are homogeneous circumscribed tumors (oval or circular), with an echogenic bright density (evaluation with high-resolution ultrasound), and in most of cases are multiple, variable in size and site, occurring in the ventricular free wall, the ventricular septum or the atrial free walls. **Results.** Occasionally, they can become obstructive, if they become very large, and can lead to fetal hydrops, arrhyth-

mias or fetal demise. The interference with coronary arterial system may cause spontaneous death as well. Usually, rhabdomyomas are detected between 20-30 weeks of gestation. The main concern following diagnosis is the strong association of multiple rhabdomyomas with genetic conditions such as tuberous sclerosis (genetic disorder with autosomal dominant inheritance). When tuberous sclerosis is diagnosed clinically after birth, in 50-80% rate of cases it is associated with rhabdomyomas. **Conclusions.** Rhabdomyomas are the main cardiac tumor, detected between 20 and 30 weeks of gestation, in 40% of cases existing fetal intracranial lesions. It is commonly associated with tuberous sclerosis, and now the diagnosis is achieved by molecular genetic testing.

Keywords: rhabdomyoma, cardiac tumors, tuberous sclerosis

Epidemiological aspects of SARS-CoV-2 infection in the Obstetrics-Gynecology Clinic of Sibiu

Radu Chicea

Faculty of Medicine, "Lucian Blaga" University of Sibiu, Romania

Introduction. Coronavirus disease is a pandemic disease with a rapid increase in cases and deaths since its first identification in Wuhan, in December 2019. The aim of the study is to evaluate epidemiologic and prognostic factors for the deliveries assisted in the Obstetrics and Gynecology Clinic of Sibiu since March 2020. Frequent manifestations include fever, cough, myalgia, headache and diarrhea. Abnormal testing includes abnormalities on chest radiographic imaging, lymphopenia, leukopenia and thrombocytopenia. **Materials and method.** We have evaluated all the deliveries from pregnancies complicated with COVID-19 during since March 2020. The data were obtained from the written information in the observation papers and in the delivery registers. There were obtained general data, delivery mode, age, gestational age, fetal weight, modality of delivery, newborn evaluation at delivery and *in utero* transmission of the disease. **Results.** A total of 85 pregnancies were assisted during the period in

observation, the average age of the pregnant women was 32.4 years old, the average gestational age was 38.1 weeks, 40 deliveries being assisted in the natural way and 45 by caesarean section. The main three indications for caesarean section were scarred uterus, advanced maternal age and thrombophilia. The average weight of the newborn was 3120 grams. No case of intrauterine transmission to the fetus was found. No severe complications, neither obstetrical nor respiratory, were registered. **Conclusions.** Because coronavirus disease might increase the risk for pregnancy complications, the management should optimally be in a healthcare facility with close maternal and fetal monitoring. No evidence of *in utero* transmission was seen in severe acute respiratory syndrome. Currently, no coronavirus-specific treatments have been approved by the Drug Administration in Europe or in the United States of America.

Keywords: COVID-19, epidemiology, obstetrics

The challenges of diagnosis and management of pregnancy associated with neurofibromatosis and Becker nevus syndrome. Report of a unique clinical case

Monica Mihaela Cîrstoiu¹, Natalia Țurcan², Anca Simionescu³, Octavian Munteanu², Delia Grădinaru-Fometescu², Florina Nedelea⁴

1. Department of Obstetrics and Gynecology, "Carol Davila" University of Medicine and Pharmacy; Bucharest University Emergency Hospital, Romania

2. Department of Obstetrics and Gynecology, Bucharest University Emergency Hospital, Romania

3. Department of Obstetrics and Gynecology, "Carol Davila" University of Medicine and Pharmacy, Bucharest; "Filantropia" Clinical Hospital, Bucharest, Romania

4. Department of Medical Genetics, "Filantropia" Clinical Hospital, Bucharest, Romania

Introduction. Neurofibromatosis is a multiorgan affecting genetic disorder with multiple implication in a pregnancy course. Most frequent pregnancy-related complications in this background are first-trimester spontaneous abortion, stillbirth, intrauterine growth restriction and a high rate of caesarean section. Becker nevus syndrome is typically characterized by the presence of circumscribed cutaneous hamartoma, hyperpigmented and hypertrichotic, along with unilateral breast hypoplasia, muscle, skin and/or skeletal abnormalities. The association between these two disorders is an extremely rare event. Moreover, pregnancy in the context of these two disorders overlapped is an exceptional entity. **Case presentation.** We present

the case of a 24-year-old primiparous gravida, 25 weeks of gestation, who was admitted to our clinic for a multidisciplinary approach in the context of severe deformation of the cervical-thoracic-lumbar spine, in "C" shape, associated with multiple café-au-lait spots, suggestive for neurofibromatosis type 1, and an asymptomatic large hyperpigmented and hypertrichotic lesion in the lumbar region, clinically indicating Becker nevus syndrome. **Conclusions.** The course of this case was clearly a challenge, being a unique case in terms of reports in the specialized literature, both nationally and internationally.

Keywords: neurofibromatosis, Becker nevus syndrome, pregnancy

Therapeutic management in advanced vulvar cancer

Bogdan Cătălin Coroleucă, Ciprian-Andrei Coroleucă, Andrei Manu, Alexandra Băușic, Diana-Elena Comandașu, Diana Mihai, Elvira Brătîlă

"Prof. Dr. Panait Sârbu" Clinical Hospital of Obstetrics and Gynecology, Bucharest; "Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

Introduction. Vulvar cancer is the fourth most common gynecological malignancy. According to predisposing factors, vulvar cancer can be categorized into two types: the first type is found in younger women and is correlated with HPV infection. The second type is independent of HPV and occurs in elderly women. Squamous cell carcinoma accounts for 90% of the vulvar cancer. Early-stage disease is defined as stage I or II. Locally advanced disease is defined as stage III or IVA. Stage IVB disease refers to women with distant metastases. **Materials and method.** We present a systematic review in which we included articles concerning the treatment of advanced vulvar cancer. We will also present several cases of advanced vulvar cancer in order to demonstrate the right management of such cases. **Results.** The treatment of vulvar cancer involves a combination of surgery, radiotherapy and, rarely, chemotherapy. If the disease is locally advanced and the lesion is ≥ 1 cm from the median line, a radical vulvectomy and ipsilateral dissection of the inguino-femoral lymph nodes are performed. If the lesion is in median line, a radical vulvectomy and bilateral dissection of the inguino-femoral lymph nodes are

performed. If the lymph nodes are positive, it can be opted for external beam radiotherapy of the primary tumor, lymph node and pelvis. If the nodes are negative, external beam radiotherapy of the primary tumor is performed. In all cases, adjuvant treatment follows. In patients with large tumors (above 4 cm) and macroscopic nodal metastases or extended localized disease to adjacent organs, the treatment course consists of chemoradiation with external beam radiotherapy to the tumor, groin and pelvis, as well as chemotherapy and surgery with perineal reconstruction. **Conclusions.** The basic treatment for vulvar cancer is surgery, but the radical nature of the procedure and adjuvant chemoradiation depends on the affectation of the lymph nodes in the inguino-femoral region and on the size of the tumor. The treatment can be long lasting and have a major impact on the quality of life. Perineal reconstruction can be accomplished using vulvoplasty, musculocutaneous flap, gracilis muscle flap, local flaps (transposition flaps) and biological graft flaps, with good results.

Keywords: vulvar cancer, surgery, radical vulvectomy, chemoradiation therapy, inguinal lymphadenectomy

Laparoscopic hysterectomy in Bega clinic from Timișoara

Marius Craina, Elena Bernad, Carmen Silaghi, Adelina Erimescu, Zoran Popa, Radu Neamțu, George Dahma

Department of Obstetrics and Gynecology, "Victor Babeș" University of Medicine and Pharmacy, Timișoara, Romania

Introduction. The patient with uterine pathology currently benefits from minimally invasive treatment, which brings essential advantages compared to open surgery. Laparoscopic hysterectomy is the ideal solution for patients who have indication for this type of approach. This paper aims to highlight the incidence and evolution of the cases treated in Bega Clinic, Timișoara.

Materials and method. A retrospective study was conducted between 2017 and 2019 in the obstetrics-gynecology clinics of the "Pius Brînzeu" County Emergency Clinical Hospital from Timișoara. The inclusion criteria were: the patient who underwent a total laparoscopic hysterectomy. In addition, the pathologies for which the treatment was applied, other associated pathologies if identified, the used surgical therapy, the evolution and

the complications were evaluated. **Results.** Most indications for laparoscopic hysterectomy were the benign pathology, which was confirmed by histopathological examination of the removed specimens. Preoperative evaluation of the case is critical to identify potential risk factors for laparoscopic surgery on the patient. The evolution of the circumstances and the complications identified both intraoperative and postoperative were noted. The importance of team cooperation was highlighted to obtain better communication and coordination, so it would be possible to improve the results in these cases. **Conclusions.** When it comes to performing a hysterectomy, laparoscopic surgery is a preferred method by both physicians and patients, due to multiple benefits.

Keywords: laparoscopy, hysterectomy, fibroma

The value of laparoscopic surgery in adnexal torsion

Marius Craina, Elena Bernad, Carmen Silaghi, Adelina Erimescu, Zoran Popa, Radu Neamțu, George Dahma

Department of Obstetrics and Gynecology, "Victor Babeș" University of Medicine and Pharmacy, Timișoara, Romania

Introduction. Adnexal pathology can be one of the causes of acute abdomen, especially in cases where torsion of the adnexa occurs. These cases usually represent a surgical emergency and sometimes require an immediate decision on treating the patient. This paper aims to identify the frequency of this pathology and how to treat it in Bega Clinic, Timișoara. **Materials and method.** Patients who benefited from laparoscopic surgical treatment for pelvic/abdominal pain syndrome that was proven intraoperatively to be caused by the torsion of the adnexa were included in this retrospective study. The study was conducted in the obstetrics-gynecology clinics of the "Pius Brînzeu" County Emergency Clinical Hospital from Timișoara, between 2017 and 2019. Both

preoperative and intraoperative aspects, and also the histopathological results, were noted. **Results.** Adnexal torsion was diagnosed in 16 cases. Eleven cases were solved laparoscopically. Both pathological aspects of the ovaries and the fallopian tubes were present, which were the cause of the torsion of the adnexa. The evolution of the cases and the intraoperative and postoperative complications were noted. **Conclusions.** Ultrasonography remains a valuable investigation to which every gynecologist can have access if they suspect an adnexal torsion. However, immediate laparoscopic surgery is currently the ideal solution for these cases.

Keywords: adnexal torsion, acute abdomen, laparoscopic surgery

Preeclampsia – screening, monitoring and clinical cases

Oana Eliza Crețu^{1,2}, Alina-Alexandra Dîrlău^{1,2,3}, Cristian Poalelungi^{1,2}, Dragoș Dobrițoiu^{1,2}, Luliana Ceaușu^{1,2}

1. "Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

2. Obstetrics and Gynecology Department, "Dr. Ioan Cantacuzino" Clinical Hospital, Bucharest, Romania

3. "Dr. N. Kretzulescu" Medical Center for Diagnosis and Treatment, Bucharest, Romania

Introduction. Preeclampsia is one of the top five causes of maternal and perinatal morbidity and mortality worldwide and is associated with a higher risk of complications for both mothers and babies. Great advances in preeclampsia screening have been made over the decades. The key to maintaining optimal screening performance is to establish standardized protocols for biomarker measurements and regular biomarker quality assessment. Traditional first-trimester screening, which is based on maternal factors and biomarkers, is the best way to identify the high-risk group and to initiate therapeutic intervention, reducing the rate of disorder complications. Very close maternal-fetal surveillance is required in management of preeclampsia. **Method.** We present the management of four early-onset preeclampsia cases followed-up in the "Dr. I. Cantacuzino" Clinical Hospital, Bucharest. The first- and second-trimester screening revealed an increased risk for developing preeclampsia before 34 weeks of gestation. Therefore, timely therapeutic intervention and careful monitoring were required. **Results.** Two of the patients developed

preeclampsia at 28 weeks of gestation, one at 29 weeks of gestation and the fourth one at 22 weeks of gestation. Preterm delivery was required due to maternal and fetal complications. Screening of the patients and appropriate monitoring lead to successful management of the cases.

Discussion. A major challenge in modern obstetrics is the early identification of pregnancies at high risk of early onset preeclampsia and undertaking the necessary measurement to improve placentation and reduce the prevalence of the disease. Various first-trimester prediction models have been developed. The American College of Obstetricians and Gynecologists (ACOG) and the National Institute for Health and Care Excellence (NICE) have proposed a screening based on maternal risk factors. The Fetal Medicine Foundation first-trimester prediction model, which includes a combination of maternal factors, measurements of mean arterial pressure, uterine pulsatility index and placental growth factor, has higher detection rate.

Keywords: preeclampsia, screening for preeclampsia, guidelines

Implications of hypertriglyceridemia-triggered acute pancreatitis during pregnancy

Gheorghe Cruciat^{1,2}, Georgiana Nemeti², Iulian Goidescu^{1,2}, Andreea Florian²

1. Emergency Hospital, Cluj-Napoca County Clinical Hospital, Romania

2. "Iuliu Hatieganu" University of Medicine and Pharmacy, Cluj-Napoca, Romania

Acute pancreatitis (AP) is a rare event during pregnancy, but its occurrence poses both mother and fetus at substantial risk. AP peaks in the third trimester of pregnancy or early during the postpartum period. Among the etiologies of acute pancreatitis, hypertriglyceridemia triggered disease must be suspected in high-risk patients. Triglyceride levels exceeding 1000 mg/dL are considered the threshold for suspicion of severe pancreatitis onset. There are no pregnancy-specific diagnostic and

therapeutic protocols. Close follow-up and prophylactic treatment must be applied to high-risk patients. Scarce pharmacologic options are amenable during pregnancy for diagnosed hypertriglyceridemia-triggered acute pancreatitis, while the use of apheresis is still under debate. The obstetric context may complicate decision-making and bears potential legal implications.

Keywords: hypertriglyceridemia, acute pancreatitis, pregnancy

Comparison of demographic and ultrasound regression formulas for hypertensive pregnancy complications

Voicu Dașcău¹, Gheorghe Furău¹, Luminița Pilat², Cristina Onel¹, Adeline Horga³, Iman Ait El Haj¹, Cristina Popescu⁴, Maria Pușchița⁵

1. Department of Obstetrics and Gynecology, "Vasile Goldiș" Western University, Arad, Romania

2. Department of Biochemistry, "Vasile Goldiș" Western University, Arad, Romania

3. Department of Pediatrics, "Vasile Goldiș" Western University, Arad, Romania

4. Department of Life Sciences, "Vasile Goldiș" Western University, Arad, Romania

5. Department of Internal Medicine, "Vasile Goldiș" Western University, Arad, Romania

Introduction. Hypertensive complications of pregnancy can often lead to serious, even life-threatening situations for the mother and fetus. First-trimester uterine artery Doppler can be a predictive tool for such complications. **Materials and method.** We have examined 305 pregnant patients from 11 weeks to 13 weeks+6 days gestational age by ultrasound in our hospital, both as inpatients and outpatients, and we have assessed the evolution of their pregnancy. We have calculated the values of AUC for hypertensive pregnancy complications for different combinations of demographic and/or ultrasound factors. **Results.** There were 21 patients with hypertensive pregnancy complications (6.89%), and 284 with normal outcome (93.11%). Three patients had gestational

hypertension (GH; 0.98%), and 18 had preeclampsia (PE; 5.90%), with 12 cases of mild PE (3.95%) and six cases of severe PE or 1.97%). The AUC values ranged from 0.510 to 1.000 for GH, 0.507 to 0.934 for mild PE, 0.505 to 0.944 for severe PE, 0.532 to 0.870 for all PE, and 0.506 to 0.885 for all hypertensive complications. The range of the mean values of each separate combination was 0.577-0.926. GH had the highest mean of AUC values, with 0.767. The total mean value was 0.690. **Conclusions.** The combination of all available ultrasound and demographic factors offers the highest AUC values for all hypertensive pregnancy complications in our study.

Keywords: screening, pregnancy, uterine artery, preeclampsia, area under curve

Statistical comparison of regression formulas for hypertensive pregnancy complications

V. Dașcău¹, Gheorghe Furău¹, Luminița Pilat², Cristina Onel¹, Adeline Horga³, Iman Ait El Haj¹, Cristina Popescu⁴, Maria Puschita⁵

1. Department of Obstetrics and Gynecology, "Vasile Goldiș" Western University, Arad, Romania

2. Department of Biochemistry, "Vasile Goldiș" Western University, Arad, Romania

3. Department of Pediatrics, "Vasile Goldiș" Western University, Arad, Romania

4. Department of Life Sciences, "Vasile Goldiș" Western University, Arad, Romania

5. Department of Internal Medicine, "Vasile Goldiș" Western University, Arad, Romania

Introduction. Hypertensive complications of pregnancy can often lead to serious, even life-threatening situations for the mother and fetus. First-trimester uterine artery Doppler can be a predictive tool for such complications. **Materials and method.** We have examined 305 pregnant patients from 11 weeks to 13 weeks+6 days gestational age by ultrasound in our hospital, both as in and outpatients, and we have assessed the evolution of their pregnancy. We have calculated the values of AUC for hypertensive pregnancy complications for different combinations of demographic and/or ultrasound factors without and with multiples of median (MoM) of pregnancy associated plasma protein-A (PAPP-A) and we have compared the results of the two studies. **Results.** There were 21 patients with hypertensive pregnancy complications (6.89%), and 284 with normal outcome (93.11%). Three patients had gestational hypertension

(GH; 0.98%), and 18 had preeclampsia (PE; 5.90%), with 12 cases of mild PE (3.95%) and six cases of severe PE (1.97%). The study including PAPP-A offered the majority of higher AUC values for GH (61.11%), severe PE (88.89%), all PE (75%) and all hypertensive complications (52.78%) and the higher mean AUC value for mild, severe and all PE, while the ultrasound study offered the majority for mild PE (58.33%) and the higher mean AUC value for GH and all hypertensive complication. The only statistically significant difference was that between the mean AUC values of the two studies for severe PE ($p=0.0004$). **Conclusions.** The combination of all available ultrasound and demographic factors offers the highest AUC values for all hypertensive pregnancy complications in our study.

Keywords: screening, pregnancy, uterine artery, preeclampsia, area under curve

The benefits of HPV vaccination beyond prophylaxis

D. Diculescu, D. Mihiu, C. Iuhaș, Renata Nicula, Carmen Bucuri, A. Mălușan, R. Ciortea, Marina Dudea, Maria Rada, Daria Pop

Second Department of Obstetrics-Gynecology, County Emergency Clinic Hospital Cluj-Napoca; "Iuliu Hațieganu" University of Medicine and Pharmacy, Cluj-Napoca, Romania

Vaccination against human papillomavirus (HPV) was introduced into clinical practice as the most effective strategy for primary prevention of cervical cancer and HPV-induced benign or premalignant lesions. It has been demonstrated that not only adolescents but also adults respond positively to vaccine immune stimulation. A history of abnormal Pap test and genital warts or HPV infection may reduce the efficacy of vaccination, but does not contraindicate it. HPV vaccination increases the magnitude and quality of natural immunity. Radical

treatments of cervical lesions do not ensure the elimination of HPV infection. Using the vaccine without the surgical removal of the HPV lesion is useless. However, combining surgical treatment and HPV vaccination can play a crucial role in reducing recurrences. Evidence supports the fact that universal vaccination, of both men and women, from adolescence to 45-50 years old, would represent the optimal modality to control HPV infection-related diseases.

Keywords: HPV vaccination, prophylaxis

Thrombophilia-related and placenta-mediated pregnancy complications

Flavia Andreea Dincă¹, Marinela Ionela Raicea¹, Elena Loredana Ciurea¹, Oana Grigorie², Diana Mitran², Costin Berceanu³

1. Department of Obstetrics and Gynecology, County Emergency Clinical Hospital, Craiova, Romania

2. Doctoral School of Medicine, University of Medicine and Pharmacy of Craiova, Romania

3. Department of Obstetrics and Gynecology, County Emergency Clinical Hospital Craiova; University of Medicine and Pharmacy of Craiova, Romania

Introduction. The aim of this paper is the integrated approach on the influence of thrombophilia on pregnancy and trophoblastic or placental structure, to study the impact of clinical conditions associated with maternal thrombophilia on the placenta, and their correlation with ultrasonographic (US), morphological and immunohistochemical study of placental angioarchitecture. **Materials and method.** This is a retrospective and prospective study, observational and descriptive, accomplished in five years (2017-2021), with a total of 473 selected pregnant women diagnosed with thrombophilia (TG), and a number of 317 nonthrombophilic pregnant women (nTG) with or without placenta-mediated pregnancy complications (PMPC). All patients included in the study were thrombophilia tested. The patients included in the study group were Caucasian, with a mean age of 30 years old (17-43). Following the application of the inclusion/exclusion criteria, 439 patients were retained in the TG group. **Results.** The tests performed showed that, out of a total of 439 cases (TG), 357 (81.32%) represented hereditary thrombophilias, 79 (17.99%) of them were acquired thrombophilias, and three cases (0.68%) had an equivocal result. The macroscopic analysis of the placenta and umbilical cords showed that most morphological changes are more common in the TG group, thus suggesting the involvement of thrombophilia in the patho-

genesis of PMPC. Placental histological and immunohistochemical changes are configured in a spectrum of nine microscopic findings. Obstetric thrombophilia specific therapy included, under various treatment regimens, acetylsalicylic acid, low-molecular-weight heparins, folic acid and supplements containing vitamins B6 and B12. **Conclusions.** Thrombophilia impacts the placental structure, especially in its angioarchitecture. PMPC occur with a relatively higher incidence in thrombophilia than in nonthrombophilic cases. Obstetric thrombophilias are associated with an increased risk of perinatal complications. The US anomalies of the fetal and maternal-fetal Doppler hemodynamic profile have as background the morphological changes that imply the alteration of the placental angioarchitecture. By integratively analyzing the clinical observations, US Doppler, but especially the morphological and immunohistochemical ones, a placental morphological pattern of thrombophilic type can be outlined. Antithrombotic or anticoagulant therapy should be accurately administered, either preconceptionally or from the onset of pregnancy, and then continued to be differentiated, until the second trimester or even until the end of pregnancy, aspects dictated primarily by the diagnosis of the thrombophilia type.

Keywords: thrombophilia, placenta, morphology, immunohistochemistry, angioarchitecture

COVID-19 pandemic effects on maternal follow-up and anemia as a pregnancy complication – experience from a tertiary care center

Alina-Alexandra Dîrlău^{1,2,3}, Oana-Eliza Crețu^{1,2}, Delia-Cristina Angelescu^{1,2}, Alice Ionescu^{1,2}, Cristina-Ioana Iancu^{1,2}, Djamila Djafer¹, Cristian Poalelungi^{1,2}, Dragoș Dobrițoiu^{1,2}, Iuliana Ceaușu^{1,2}

1. "Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

2. "Dr. Ioan Cantacuzino" Clinical Hospital, Bucharest, Romania

3. "Dr. Nicolae Kretzulescu" Medical Centre for Diagnosis and Treatment, Bucharest, Romania

Introduction. The current outbreak of coronavirus (COVID-19) disease, declared by WHO as a global pandemic on 11 March 2020, comes with unprecedented effects on health systems worldwide, pregnant women and their fetuses being considered a high-risk population. **Objectives.** Our study aims to evaluate the impact of the COVID-19 pandemic on pregnancy and peripartum anemia, one of the most frequent and important complications of pregnancy, with severe effects on maternal and neonatal health. **Materials and method.** A retrospective observational study was conducted in the Obstetrics and Gynecology Department of the "Dr. I. Cantacuzino" Clinical Hospital, a tertiary care maternity in Bucharest. The study included 1824 pregnant women with anemia, out of 2962 births that were admitted in our hospital between the 1st of April 2019 and the 31st of March 2021, comparing maternal and neonatal outcomes one year before and one year after the COVID-19 outbreak, respecting current diagnostic criteria recom-

mended by WHO and regional guidelines (hemoglobin concentration <11 g/dL before/after birth). **Results.** The results showed that the pandemic group had poor or no follow-up during the pregnancy compared with the period before, had more severe anemia, needed more intravenous iron therapy or blood products administration, with the necessity of longer hospitalization periods for the mothers and the neonates and higher peripartum complications and premature birth rate. **Conclusions.** Anemia is one of the most frequent complications of pregnancy and birth, which, in our experience, was worse in the global context of COVID-19 outbreak, mostly due to insufficient prenatal care. Future research needs to focus on pregnancy anemia impact for maternal and infant health and on developing strategies of increasing the follow-up surveillance to reduce anemia among pregnant women.

Keywords: maternal anemia, iron depletion, pregnancy complications, COVID-19 and pregnancy

Why use an ultrasound evaluation in the labor ward?

Roxana-Cristina Drăgușin¹, Laurențiu Dîră², Maria Șorop-Florea³, Marina Dinu¹, Lucian Zorilă¹, Rodica Nagy⁴, Alina Enache⁴, Anca-Maria Istrate-Ofițeru⁵, Dominic-Gabriel Iliescu¹

1. Department of Obstetrics and Gynecology, University of Medicine and Pharmacy of Craiova; University Emergency County Hospital of Craiova, Romania

2. "Filantropia" Clinical Hospital, Craiova, Romania

3. Department of Obstetrics and Gynecology, University of Medicine and Pharmacy of Craiova; "Filantropia" Clinical Hospital, Craiova, Romania

4. University Emergency County Hospital, Craiova, Romania

5. Department of Histology, University of Medicine and Pharmacy of Craiova; University Emergency County Hospital, Craiova, Romania

Introduction. In obstetric practice, every physician is aiming to shun from any difficult vaginal delivery, as it can lead to serious fetal and maternal complications. Both doctor and pregnant patients at term who are admitted in the labor ward want to know the chance for a successful vaginal delivery and the risks for an operative delivery (instrumental delivery or caesarean section). Clinical evaluation remains the gold standard for managing labor. However, the use of intrapartum ultrasound (IPU) has been proposed to help physicians in managing labor as a noninvasive method to predict obstructed labor in low-risk patients and to identify high-risk patients even before the onset of the partum. **Method.** We reviewed all studies demonstrating that ultrasound examination is more accurate and reproducible than the clinical

examination for evaluating cervical dilatation, fetal head station, and fetal head position. **Results.** Discovering and using a risk tool may be useful for reassuring most women regarding their likelihood success at achieving an uncomplicated vaginal delivery, as well as selecting those patients with such a high risk for caesarean delivery that they should avoid a trial of labor. Such a risk tool has the potential to highly improve planning hospital service needs and minimizing the patients' risk. **Conclusions.** In the last decade, IPU assessment has become a need, as it proved to be a reliable and objective method for obstetricians in decision making and preventing unnecessary intervention. IPU should be used as a complementary tool in labor and not undermine the clinician's skills.

Keywords: intrapartum, ultrasound, prolonged labor

Vulvar cancer – a case report of a patient who presented vulvar squamous cell carcinoma with rapid disease progression

Mihai Cristian Dumitrașcu^{1,2}, Florica Șandru^{2,3}, Olga Scaletchi¹, Răzvan Fodoriu¹

1. Bucharest University Emergency Hospital, Romania

2. "Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

3. Department of Obstetrics-Gynecology and Neonatology, "Elias" University Emergency Hospital, Bucharest, Romania

Vulvar cancer is a rare malignancy of female genitalia that involves vaginal lips, the opening of the vagina and the clitoris. Every year, 27,000 women worldwide are diagnosed with vulvar cancer, the highest incidence being in Europe, North and South America. The pathogenesis of the disease arises from either human papillomavirus (HPV), affecting primarily younger women, or from vulvar dermatoses, such as lichen sclerosis, affecting older patients. The clinical presentation in most patients involves vulvar pruritus or vulvar lesions that present ulcerations, bleeding, pain, difficulty in urination and pain at sexual intercourse. During the COVID-19 pandemic, the access to medical care was severely restricted so, as a result, patients come in the clinics with advanced stages

of diseases. The aim of this paper is to present the case of a 53-year-old woman who was referred to our clinic for a vulvar pain and pruritus for approximately three months. The clinical examination revealed a large vulvar tumor that involved both labia majora and minora, the vaginal introitus, the clitoris and the mont pubis. Enlarged lymph nodes were palpated in bilateral inguinal region. After further investigations, we found out that the patient was biopsied a year earlier and the anatomopathological result showed a LSIL vulvar lesion with a starting point at the level of a papilloma. Unfortunately, the patient experienced fast general health deterioration, and shortly after death occurred due to generalized sepsis.

Keywords: vulvar cancer, squamous cell carcinoma

The importance of the three-vessel trachea view in prenatal screening

Roxana Gireadă¹, Răzvan Socolov¹, Elena Mihălceanu¹, Luiza Guzgă²

1. "Grigore T. Popa" University of Medicine and Pharmacy, Iași, Romania

2. "Filantropia" Clinical Hospital of Obstetrics and Gynecology, Bucharest, Romania

Introduction/aim. It is strongly recommended to include the three-vessel trachea (3VT) view in the prenatal screening scans, as the last step of a continuous cardiac sweep starting from the upper abdomen. The aim of this study was to evaluate the additional role of the 3VT in identifying congenital heart disease, compared to using the 4C view only. **Materials and method.** We reviewed the scans of 1327 low-risk pregnancies (including 10 sets of twins, of which seven bichorionic and three monochorionic) presenting for prenatal screening in a private setting. The scans were performed in the second and/or third trimester, the cardiac screening consisting of a sweep from the upper abdomen, 4C, outflow tracts, 3VT, to supraaortic area (subclavian artery and brachiocephalic vein), followed by sagittal scanning of the aortic arch and bicaval view, including Color Doppler for each view. All cardiac anomalies were extracted and then we analyzed the performance of using 4C only and 3VT

only in cardiac screening. **Results.** Twenty-eight cardiac anomalies were identified (2.1%), in the singleton pregnancies only. None associated genetic anomalies, nor were they syndromic. Using only 4C, we could identify 42.8% of anomalies, and by using only 3VT/supraaortic area we identified 71.4% of anomalies. By including 3VT/supraaortic area, 57.1% of additional anomalies were identified, 17.8% being clinically significant and also detectable in 2D mode (one dextrotransposition of the great arteries, three hypoplastic aortic arch, one pulmonary stenosis, one right aortic arch with vascular). 35.7% were isolated anatomical variants (seven aberrant right subclavian artery, three persistent left superior vena cava). **Discussion and conclusions.** The 3VT view improves the detection rate of clinically significant congenital heart disease, even without using Color Doppler.

Keywords: prenatal screening, three-vessel trachea view, congenital heart disease

Advanced breast carcinoma diagnosed during pregnancy

Iulian Goidescu, Georgiana Nemeti, Andreia Preda, Adelina Staicu, Gabriela Caracostea, Mihai Surcel, Gheorghe Cruciat, Daniel Mureșan

First Clinic of Obstetrics and Gynecology, Emergency Clinic County Hospital, Cluj-Napoca; "Iuliu Hațieganu" University of Medicine and Pharmacy, Cluj-Napoca, Romania

Introduction. Breast cancer is the most common type of cancer in women and the second leading cause of death from neoplastic disease. In recent years, there has been a decrease in the age of onset of this pathology, more women being diagnosed with breast cancer during pregnancy. Moreover, the decrease in accessibility to medical services due to the COVID 19 pandemic has led to the appearance of advanced cases of breast cancer diagnosed during pregnancy. **Case report.** We present two cases of advanced breast cancer

diagnosed in pregnancy, both patients presenting in our service for bone pain and being diagnosed with bone metastases with unknown cancer origin, which was later confirmed as breast cancer. **Conclusions.** Although pregnancy-associated breast cancer has become a more common pathology compared to previous decades, the fear of addressing medical services due to the COVID-19 pandemic leads to a late diagnosis with poor outcome.

Keywords: breast cancer, pregnancy, metastasis

Immunomodulation in genital HPV infections – in between mirage and real benefit

Corina Grigoriu^{1,2}, Gina Ionescu², Andra Bălan², Irina Horhoianu^{1,2}

1. Department of Obstetrics and Gynecology, "Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

2. Bucharest University Emergency Hospital, Romania

Introduction. HPV is nowadays a frequent infection after sexual life commencement. Due to affordable information sources, patients begin to understand the interconnection between the aggressivity of the viral infection and the local and general defending mechanisms. Clinicians, on the other hand, are also interested in resolving as quick as possible these annoying infections. **Materials and method.** Questions we try to answer are: Is there a real benefit in some enhancing of the natural host defending mechanisms? Is coping with stress important even for an HPV-infected patient? Does smoking play a real part in this game? What about vaginal microbioma? As far as we know at this moment, patients with alterations of the vaginal microbioma are more prone to have persistent HPV infections and, as such, are at higher risk for developing high grade cervical intraepithelial neoplasia. But why do these imbalances in vaginal flora occur? **Results.** Beginning with advices toward a healthier lifestyle, to stopping smoking and

a better stress control, most patients expect from the clinician some sort of miraculous treatment in order to enhance the defending mechanisms against HPV genital infection. Most treatments are over-the-counter available and information is overflowing the internet sites and forums. The question is: do they really work, for how long, at what costs and with what benefit? In use there are local treatments (immunomodulators – natural or synthesis, ovules, sprays, creams) or general immunomodulators (isoprinosine, orthomolecular supplements). Alternative therapies (herbal therapies, homeopathy, acupuncture, Chinese medicine) are reported to have some results. Therapeutic vaccines and specific immunotherapy are intensively studied. **Conclusions.** Immunomodulation in genital HPV infection is still an area of quicksand and personalized medicine is for the moment the answer in this domain.

Keywords: general and local immunomodulation, genital HPV

Large loop excision of the transformation zone: “see and treat” or “think twice about it”?

Corina Grigoriu^{1,2}, Gina Ionescu², Andra Bălan², Irina Horhoianu^{1,2}

1. Department of Obstetrics and Gynecology, “Carol Davila” University of Medicine and Pharmacy, Bucharest, Romania

2. Bucharest University Emergency Hospital, Romania

Introduction. Large loop excision of the transformation zone (LLETZ) is a valuable procedure for the management of cervical intraepithelial dysplasia. Several advantages are certain: it is an outpatient procedure, it may be both diagnostic and therapeutic, the cervical tissue heals rapidly, the future obstetrical outcome is satisfactory. **Materials and method.** But all these advantages conduct sometimes to a hasty attitude – the “see and treat” approach, performing the procedure immediately after a colposcopy, or even worse, immediately after receiving a slightly modified cervical cytological result (without even visualizing the source-tissue through the colposcope). The “see and treat” approach is justified in thoroughly selected cases: absence of any cervical infection, absence of vaginosis, postmenstrual period, visualization of the entire transformation zone

(which is obviously possible only during colposcopy), the colposcopic impression of moderate or high-grade dysplasia; the patient lives remote from the office or may be lost for follow-up. **Results.** Maybe sometimes it’s better to “think twice about LLETZ”. The consequences of an unnecessary rush may be moderate, but also severe: chronic cervicitis, chronic pelvic pain, deep dyspareunia, repeated and worrisome episodes of vaginal bleeding, infertility, risk of abortion or premature birth. Postponing the procedure for no more than one month in selected cases may be more advantageous for the patient than a life-time burden in gynecological pathology. **Conclusion.** The precept *Primum non nocere, deinde curare* has to be respected in the management of cervical dysplasia.

Keywords: LLETZ, indications, risks

Role of pulse oximetry in the screening of congenital heart diseases

Monica Gabriela Hășmășanu¹, Liliana Kalisch², Margareta Părvănoiu², Melinda Matyas¹, Ligia Blaga¹, Gabriela Zaharie¹

1. Department of Neonatology, “Iuliu Hațieganu” University of Medicine and Pharmacy, Cluj-Napoca, Romania

2. Department of Neonatology, “Dr. Constantin Opreș” County Emergency Hospital, Baia Mare, Romania

Introduction. Congenital heart defects are the most common malformations and a quarter of them are critical and often require surgery or catheterization in the first weeks of life. The early detection of them is essential for a good outcome. **Materials and method.** This is a prospective descriptive study, performed over a period of three months, on a number of 550 newborns, born in level 2 maternity. We analyzed a group of 541 newborns hospitalized in the Neonatology Department of the Clinical Hospital Baia Mare, in which pulse oximetry screening was performed, and in those with positive screening, cardiac ultrasound was performed. On the third day of life, pulse oximetry screening was performed, data were collected on the mother and pregnancy and on the newborn, respectively. **Results.** We analyzed 541 births, of which 318 (58.78%) natural births and 223 (41.22%) births by caesarean section. We obtained positive screening in 79 cases. The characteristics of the group with positive screening are: average weight

3068.41±680 grams; average gestational age 37.5±2.75 weeks, preductal saturation of 92%±0.04, and postductal saturation 90%±0.04. According to the screening, there is a higher percentage of saturations between 91% and 95% (57.26%), compared to O₂ saturations that have a difference of more than 3% between pre- and postductal saturation, the percentage being 42.73%. Analyzing resuscitation at the delivery room, we have 41.77% needed free-flowing O₂, 20.25% of newborns needed CPAP-type ventilator support, and 1.26% needed intubation. Of the cases with positive screening, in eight cases echocardiography confirmed: interventricular septal defect (four cases), interatrial septal defect (three cases) and pulmonary stenosis (one case). **Conclusions.** Congenital heart disease was confirmed in 10.12% of newborns with positive screening. Pulse oximetry screening is a simple, noninvasive, painless and inexpensive way to diagnose congenital heart defects.

Keywords: congenital heart, newborn, screening

The late preterm infants – a continuous challenge

Monica Gabriela Hășmășanu¹, Bianca Marcela Voina², Melinda Matyas¹, Ligia Blaga¹, Dan Boitor³, Andrada Mădălina Petyan², Gabriela Zaharie¹

1. Department of Neonatology, "Iuliu Hațieganu" University of Medicine and Pharmacy, Cluj-Napoca, Romania

2. "Iuliu Hațieganu" University of Medicine and Pharmacy, Cluj-Napoca, Romania

3. First Department of Obstetrics and Gynecology, "Iuliu Hațieganu" University of Medicine and Pharmacy, Cluj-Napoca, Romania

Introduction. The late preterm is the newborn of the gestational age of 34-36 weeks with 6 days, also called near term or moderately premature. Although considerable progress has been made in gynecology and neonatology in recent years, prematurity has remained the important cause of neonatal morbidity and mortality. **Materials and method.** This paper is a descriptive study, at 12 months, performed by the Neonatal Intensive Care Unit (NICU) within the Obstetrics-Gynecology I, Cluj-Napoca. We analyzed data on pregnancy, mother and neonatal data. We evaluated the cost within the two compartments: NICU and Premature. **Results.** The study group included 142 newborns with late preterm born in the clinic or transferred from level 1 or 2 units. In the study group, female prematurity predominated. The gestational age of 36 weeks predominated in the study group, as well as the weight between 2000 and 2490 grams. Caesarean section predominated as a way of birth. Analyzing pregnancy pathology in the study group, pregnancy-induced hypertension and ges-

tational diabetes predominated. Respiratory pathology was present in 87% of cases due to respiratory distress caused by surfactant deficiency, transient tachypnea and pneumothorax. 30% of them required noninvasive respiratory support with an oxygen demand in the first day of life of 28.58±11.68%. During the hospitalization, in 82% of cases hyperbilirubinemia was encountered, and hypoglycemia in 8%. The expenses during the hospitalization are 2993.57±213.3 RON on the NICU department and 1787.67±89.2 RON on the premature department. **Conclusions.** Late prematurity is an important factor in hospitalization in the neonatal intensive care unit. The main risk factors for late prematurity are pregnancy-induced hypertension and gestational diabetes. Early morbidities in the newborn with late prematurity were respiratory distress due to surfactant deficiency and hyperbilirubinemia. Late premature involves significant costs in both the intensive care unit and the premature department.

Keywords: prematurity, late preterm, morbidity

Inherited thrombophilia in pregnancy: a cause of postpartum hemorrhage?

Ciprian Ilea^{1,2}, Demetra Socolov^{1,2}, Irina Stoian¹

1. Department of Obstetrics and Gynecology, Faculty of Medicine, "Grigore T. Popa" University of Medicine and Pharmacy, Iași, Romania

2. "Cuza Vodă" University Hospital, Iași, Romania

Objective. Postpartum hemorrhage (PPH) is the leading direct cause of maternal death and most difficult to control are hemorrhages associated with coagulation disorders. The aim of this study was to determine whether antithrombotic therapies increase the risk of PPH and puerperal genital hematomas. **Method.** The medical records of patients with hereditary thrombophilia and PPH or puerperal genital hematomas were reviewed retrospectively. The following baseline and obstetric data were collected: age, gravidity, parity, body mass index, duration of pregnancy, history of previous deep vein thrombosis or adverse pregnancy outcome, type of birth, onset of postpartum hemorrhage and maternal hematomas, and need for transfusion of blood products. In case of vaginal

births, the following were considered: type of labor and delivery, duration of labor, and birth weight. **Results.** In only 12 cases, we found a possible direct association between the use of thromboprophylaxis with low-molecular-weight heparin (LMWH) during pregnancy and the occurrence of PPH or puerperal hematomas (vaginal/vulvar, broad ligament, uterovesical fold, and epidural after spinal anesthesia). **Conclusions.** The need to evaluate the PPH in women receiving heparin prophylaxis became obvious in pregnancies complicated by inherited thrombophilia, although the lack of a clear association is likely attributable to the multifactorial nature of PPH.

Keywords: pregnancy, inherited thrombophilia, low-molecular-weight heparin, postpartum hemorrhage

The importance of diagnosis of uterine congenital malformations in pregnant patients

Alice Ionescu^{1,2}, Delia Angelescu^{1,2}, Alina-Alexandra Dîrlău^{1,2}, Oana-Eliza Crețu^{1,2}, Cristina Iancu^{1,2}, Alexandru Ciulcu², Iuliana Ceaușu^{1,2}

1. Department of Obstetrics and Gynecology, "Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

2. "Dr. Ion Cantacuzino" Clinical Hospital, Bucharest, Romania

Introduction. Congenital uterine abnormalities are the result of fusion and/or resorption defects of the Müllerian ducts, which appear during organogenesis. These malformations are often asymptomatic; therefore, their incidence is difficult to assess. The impact on fertility is variable and the diagnosis can be delayed by not seeing a doctor, presenting in advanced pregnancy or after complications. The most common complications are subfertility, menstrual disorders, premature birth, miscarriage, or dystocic labor, along with secondary complications from associated urinary tract malformations. **Materials and methods.** We present the case of an 18-year-old patient, gravida 2, para 2, who presented at the hospital in labor, known with scarred uterus after caesarean section (three years ago, in 2019), at a gestational age estimated at 39 weeks, with pelvic presentation, affirmatively without personal medical pathological history. The examination revealed a complete vaginal septum with separate cervixes. The patient did not present medical documents from the current or previous pregnancy. **Results.** For the prophylaxis of uterine rupture, the patient gave birth by caesarean

section to a live fetus, male, 3080 g, AI9. A septate uterus was found intraoperatively. Tubal surgical sterilization was performed upon request. The postoperative evolution was favorable. Although the importance of investigating possible associated malformations of the urinary tract was explained to the patient, she did not report to the routine check-up six weeks after giving birth. **Conclusions.** The presented case had a favorable evolution, despite the lack of monitoring and of information related to the current and previous pregnancy. Unfortunately, the vast majority of cases present with complications and the lack of data related to previous birth and/or associated pathology could be solved by better communication between specialized medical services. The early diagnosis of congenital uterine malformations may decrease the incidence of miscarriages and premature births. Also, knowing that the genital and the urinary system have a common development in the intrauterine life, it is important to investigate a possible association of malformations between the two systems.

Keywords: uterine congenital malformations, septate uterus, dystocic labor, better communication

Obstetrical soft tissue trauma during vaginal delivery in the Romanian adolescent population – a multicentric comparative study with adult population

Crîngu Ionescu¹, Alexandra Matei¹, Mihai Dimitriu¹, George Roșu¹, Dan Năvolan², Cristian Furău³

1. Department of Obstetrics and Gynecology, "Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

2. Department of Obstetrics and Gynecology, "Victor Babeș" University of Medicine and Pharmacy, Timisoara, Romania

3. Department of Obstetrics and Gynecology, Emergency County Hospital, Arad, Romania

Introduction. Romania is a country with high rates of adolescent births, associating scarce comprehensive obstetrical management in this specific population. This research aims to assess soft tissue trauma after vaginal birth in teenage mothers compared to their adult counterparts. **Materials and method.** A retrospective case-control study was conducted during one year in two hospitals. All vaginal deliveries were considered; age cut-off value was considered at 20 years old for case and control groups. Lacerations were divided into three subgroups, considering the involved anatomical region: Group I – labial and periurethral lacerations; Group II – vaginal and perineal lacerations; Group III – cervical lacerations. **Results.** There were 1498 women included in the study, 298 young mothers and 1200 adults. Teenagers were more likely to have an episiotomy during delivery compared to adult women, 56% versus 26.7%

($p=0.00$, Pearson Chi-Square). Group II lacerations were the most frequent types of birth trauma in both study groups. Fetal weight ≥ 4000 g associated a twice higher risk for vaginal and perineal lacerations when age criterion was not considered (OR=1.98; 95% CI; 1.13-3.47; $p=0.01$). Similar results were obtained in the adult reference group (OR=1.98; 95% CI; 1.08-3.62; $p=0.04$). The incidence of Group I and II lacerations increased with age, from 0% and 9.1% between 10-14 years old to 6%, and 26.2% between 18 and 19 years old. **Conclusions.** All groups of lacerations were more often identified in the case group, compared to the adult group. Fetal macrosomia could not be documented as a risk factor for obstetrical injury in young mothers.

Keywords: episiotomy, vaginal delivery, perineal trauma, soft tissue injury, birth laceration, adolescent birth

Update on maternal immunization excluding COVID-19 vaccine

Costin Alexandru Ispas¹, Antoine Edu¹, Elena Ispas², Dimitrie Nanu³

1. "Carol Davila" University of Medicine and Pharmacy, "Nicolae Malaxa" Clinical Hospital, Bucharest, Romania

2. "Elias" University Emergency Hospital, Bucharest, Romania

3. "Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

Immunization is part of medical care for both adult and children and the last year is reminding us of this field more than ever. The vaccination of the pregnant woman has proven useful not only in protecting the mother, but also the offspring in some pathologies. If influenza vaccines were not studied enough 15 years ago and were not recommended during pregnancy, today all the international medical associations consider them safe for administration at this period of a woman's life. We have reviewed the latest set of recommendations for immunization during pregnancy made by some of the major medical associations in the world, such as the American College of Obstetricians and Gynecologists

(ACOG), the European Board and College of Obstetrics and Gynecology (EBCOG), and others, and also other publications such as UpToDate, in order to assess the news in vaccination during pregnancy. The vast amount of literature emerging in the field of COVID-19 vaccination led to numerous opinions and recommendations on this issue that have already been published and disseminated in the past months, including the effects of this vaccine during pregnancy, therefore this topic was excluded.

Keywords: immunization during pregnancy, maternal immunization, maternal care during pregnancy, immunization

Induction of labor using the Foley balloon in patients with post-caesarean scar uterus

Victor Juncu^{1,2}, Camille Olivier¹, Călin Gheorghe Oancea², Mihai Emil Căpîlna²

1. Service de Médecine Foeto-Maternelle, CHU Brugmann, ULB, Bruxelles, Belgium

2. "George Emil Palade" University of Medicine, Pharmacy, Sciences and Technology of Târgu-Mureș; Emergency Clinic County Hospital, Târgu-Mureș, Romania

Introduction/objective. Patients with scarred uterus have an increased risk of uterine rupture, representing the main reason for abandon of another method to induce labor. Consequently, the scarred uterus is the most common indication for repetitive caesarean section. The aim of our paper is to evaluate the main parameters of delivery mechanism and to assess the fetal well-being at birth in patients with scarred uterus after a previous caesarean section, induced by Foley balloon, at the Brugmann University Hospital, in 2017. **Methodology.** A retrospective study has been performed for the year 2017 in a referral university hospital, analyzing the files of patients enrolled using the software Mosos-O, Mosos-CTG, QPlanner and ASTRAIA. The statistical cohort under analysis consisted of the control group (68 pregnant women with scarred uterus after a caesarean section, with the labor induced by using a Foley balloon), and the pilot group (61 pregnant women in which the labor was induced by Foley balloon for patient convenience).

Results. Patients in both study groups benefited from the onset of labor between 40 and 41 weeks of amenorrhea. Of the control group, 69% of pregnant women gave birth vaginally, compared to the pilot group in which vaginal births accounted for 70.5%. Patients from the control group (79%) had a eutocic birth compared to 63% of patients from the pilot group. The Apgar score at 5 minutes of life in the case of the control group presented in 86.86% a value higher than 7, compared to the same parameter in the pilot group (90.15%). There were no cases of uterine rupture. **Discussion and conclusions.** Labor induction using Foley balloon for pregnant women with scarred uterus after caesarean section is a feasible technique that deserves to be proposed to this profile of patients providing identical obstetric and neonatal outcome as in pregnant women with non-scarred uterus induced by the same method.

Keywords: induction of labor, post-caesarean scar uterus, Foley balloon

Considerations on COVID-19 pregnancy: a case series during 2020-2021 in Prahova region, southern Romania

Ioan Bogdan Luchian, Ali Karaouni, Călin Boeru, Letiția Goraneanu

Obstetrics and Gynecology Hospital, Ploiești

Introduction/objectives. Coronavirus 2019 (COVID-19) has rapidly spread worldwide. Previous publications mostly addressed the impact of COVID-19 on elderly population with chronic comorbidities, being at risk for severe evolution. However, data regarding the association of COVID-19 infection with pregnancy is still scarce. Therefore, we aimed to present a case series of six pregnant female patients, diagnosed with COVID-19. **Materials and method.** Six patients, with confirmed COVID-19 by real-time reverse transcription polymerase chain reaction test were admitted at the Obstetrics and Gynecology Hospital, Ploiești, Romania. Clinical information was collected on admission and during hospitalization by physicians in charge. **Results.** The mean age of the patients was 26.6 years old. Mean hospitalization duration was 11 days. The infection was diagnosed in the first 36 weeks of pregnancy. Four patients, without risk factors, presented bilateral pneumonia at 48 hours of admission. The pregnancy evolution was good, except for two cases with preterm delivery onset at 33 and 35 weeks. All four patients received standard therapy with oxygen, antibiotics and dexa-

methasone. Another patient was admitted with fever (38.5°C) and deep venous thrombosis of the left inferior limb. Anticoagulation treatment was initiated, with the complete remission of the symptoms, without impact on pregnancy. The last patient of case series is a 41-year-old female with diabetes mellitus and obesity, with a 29-week pregnancy, and admitted with high fever and severe respiratory distress syndrome. Emergency caesarean section was performed due to clinical instability. Following discharge, the patients were asymptomatic. **Conclusions.** The risk of preterm delivery might be increased by COVID-19 infection. Thus, this case series emphasizes the important role of an effective multidisciplinary approach in severe cases with high risk of complications during pregnancy, requiring continuous monitoring, prolonged oxygen therapy, anticoagulation therapy and emergency delivery. However, future studies should be conducted to evaluate the long-term evolution of patients that presented SARS-CoV-2 infection during pregnancy.

Keywords: COVID-19, pregnancy, preterm delivery, respiratory distress syndrome

Preeclampsia – a myriad of miRNAs

Andrei Mihai Măluțan^{1,2}, Doru Diculescu^{1,2}, Răzvan Ciortea^{1,2}, Cristian Iușă^{1,2}, Radu Florin Mocan-Hognogi^{1,2}, Carmen Bucuri², Maria Rada^{1,2}, Marina Ducea^{1,2}, Viorela Suciu¹, Mihai Dan^{1,2}

1. "Dominic Stanca" Obstetrics-Gynecology Clinic, County Emergency Clinical Hospital Cluj-Napoca, Romania

2. "Iuliu Hațieganu" University of Medicine and Pharmacy, Cluj-Napoca, Romania

Preeclampsia (PE), a multisystem pregnancy-associated disease, has a major contribution to maternal morbidity and mortality worldwide, affecting approximately 10 million women worldwide. PE is thought to occur as a consequence of several factors, including defective spiral artery remodeling, placental oxidative stress, endothelial dysfunction and systemic inflammation. Recently, microRNAs (miRNAs) have been associated with the pathogenesis of PE and could play an important role in the development of PE. MicroRNAs are intensely expressed in the placenta during a normal pregnancy. Thus, analyzing miRNAs in the sera of PE patients could enrich knowledge about the pathophysiological mechanisms of this disease. Recent articles highlight the relationship between the aberrant expressions of several

miRNAs and negative pregnancy outcomes. A number of miRNAs, including miR-16, miR-29b, miR-34a, miR-155, miR-210 and miR-675, have been shown to decrease the proliferation and migration of trophoblasts. At the same time, recent studies have shown that miR-210 and miR-155 are constantly aberrantly expressed in women with PE. Thus, by elucidating the functional role of these modified miRNAs, important pathways involved in PE could be identified and potential predictive/diagnostic biomarkers could be brought to light, which could be used in PE. Moreover, enriching our knowledge about the role of miRNAs in pregnancy-associated diseases is necessary in order to develop therapeutic strategies in the future.

Keywords: pregnancy, preeclampsia, miRNA

The role of adipokines in uterine contractility

Dan Mihu, Doru C. Diculescu, Răzvan Ciortea, Andrei M. Măluțan, Ciprian G. Porumb, Mihaela D. Oancea, Renata L. Nicula, Cristina M. Prodan

"Dominic Stanca" Obstetrics and Gynecology Clinic, County Emergency Clinic Hospital Cluj-Napoca, "Iuliu Hațieganu" University of Medicine and Pharmacy, Cluj-Napoca, Romania

Obesity is currently an important public health issue, registering a rapid increase in recent decades and affecting young people more and more frequently. The result is an increasing number of overweight or obese patients who reach reproductive age and become pregnant. Obesity is considered as an increased volume of adipose tissue which has well-known negative effects on multiple systems, but also on reproductive function. This piece of information led to the search for the existence of a connection between an increased Body Mass Index and the dysfunction of uterine contractility. Most cases of post-term pregnancy which usually require the induction of childbirth are observed in patients with increased Body Mass Index. The most important adipokines are represented by adiponectin, leptin, ghrelin, visfatin and apelin. Leptin has been demonstrated to act at smooth muscle fiber level in the vascular wall where it produces a decrease in calcium release from the sarcoplasmic reticulum, thus reduc-

ing the uterine contractile force. Leptin has an inhibitory effect on the frequency and amplitude of uterine contractions. Increased serum levels of leptin in obese women are involved in the inhibition of the apoptotic phenomenon involved in uterine involution. Visfatin exerts a more potent inhibitory effect compared to leptin upon uterine contractions. The inhibitory effect of ghrelin on uterine contractility involves a possible role in the regulation of myometrial activity. Apelin exerts a potent inhibitory effect on myometrial cells, causing a reduction in both the amplitude and the frequency of uterine contractions. Adipose tissue is currently regarded as a real endocrine organ, responsible for the production of bioactive polypeptides, adipocytokins. Adipokines exert modulating effects during pregnancy and there is evidence of their involvement in the pathophysiology of pregnancy-related complications.

Keywords: obesity, adipokines, uterine contractility, pregnancy, postterm

Maternal HIV infection – prematurity risk factor?

Mihai Mitran^{1,2}, Octavia Velicu^{2,3}, Elvira Brătîlă^{1,2}

1. "Prof. Dr. Panait Sîrbu" Clinical Hospital of Obstetrics and Gynecology, Bucharest, Romania

2. "Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

3. Medical Center for Outpatient Diagnosis and Treatment, Ministry of Justice, Bucharest, Romania

HIV infection represent a worldwide health issue, with 38 million known patients and around 1.7 million new cases every year. Vertical transmission from mother to child represents 2% of the new cases. Also, studies of HIV pregnancies have shown a higher rate of complications such as prematurity and intrauterine growth restriction. This paper presents a study on 78 HIV pregnancies man-

aged in the "Prof. Dr. Panait Sîrbu" Clinical Hospital of Obstetrics and Gynecology, Bucharest, between 2018 and 2020. The main conclusion of the study is that the rate of prematurity was 11.53% (9 out of 78), lower than what the literature previously reported (24%), mainly due to a very strict management of ART and pregnancy monitoring.

Keywords: HIV, pregnancy, prematurity

Magnesium sulphate as a neuroprotective agent in imminent risk for preterm delivery – a challenge for obstetrical practice

Marius Alexandru Moga, Sabina Ioana Bădilă, Oana Dimienescu, Cezar Podașcă, Cristian Arvătescu, Nicușor Bigiu, Andreea Bălan

Department of Obstetrics and Gynecology, Faculty of Medicine, Transilvania University of Brașov, Romania

Introduction/objectives. Preterm birth remains an obstetrical and neonatal challenge, despite the increasing survival rates at early gestational ages, being the leading cause of perinatal mortality and morbidity in high-income countries. Even though the survival rate depends on endogenous and exogenous factors, the risk of developing medical and neurodevelopmental impairment remains high. One of the most important adverse neurological outcomes associated with preterm birth is cerebral palsy which describes a group of permanent disorders of the development of movement and posture disorders. Magnesium sulphate is a widely available and commonly used treatment for eclampsia in most of the countries; however, its use in obstetrical practice is being controversial. The aim of this paper is to present the beneficial role and to summarize the relevant evidence and practice recommendations of magnesium sulphate in neuroprotection in infants as antenatal treatment based on the current studies. **Methodology.** We selected the studies from Google Academic and PubMed databases and reviewed

recent articles from literature, aiming to evaluate the impact of magnesium sulphate as antenatal treatment in imminent risk for preterm delivery. Our research included all the publications during the period of January 2016 to May 2021 using the following Medical Subject Headings (MeSH): “magnesium sulphate”, “neuroprotection”, “preterm delivery”, “cerebral palsy”. **Results.** Most of the studies demonstrate significant results for antenatal magnesium sulphate reducing the risk of cerebral palsy. It is unclear whether gestational age is a key factor in the neuroprotective effect of the magnesium sulphate, but several studies suggested that the outcome of death or cerebral palsy was decreased when magnesium sulphate was administered before 34 weeks of pregnancy. **Conclusions.** The main limitation of this literature review consists of the lack of long-term follow-up of the infants. Future research should focus on the dosing and timing of magnesium sulphate administration, including the factors which could predict the possible adverse reactions to this treatment.

Keywords: preterm delivery, magnesium sulphate

Pregnancy and neuromyelitis optica spectrum disorder (NMOSD) – a rare but challenging interaction

Georgiana Nemeti, Gheorghe Cruciat, Ioana Rotar, Gabriela Caracostea, Iulian Goidescu, Andreia Preda, Mihai Surcel, Dan Boitor, Daniel Mureșan

First Obstetrics and Gynecology Clinic, Mother and Child Department, “Iuliu Hațieganu” University of Medicine and Pharmacy, Cluj-Napoca, Romania

Neuromyelitis optica spectrum disorder (NMOSD) is an inflammatory central nervous system disorder affecting women of childbearing age. It is characterized by recurrent antibody-mediated inflammatory disorders targeting the optic nerve, spinal cord, area postrema of the medulla and hypothalamus. Aquaporin-4 (AQP4) is the most common target antigen in NMOSD. Disease relapses are common during pregnancy and postpartum. Younger patients, those with higher AQP4-ab titres and those with inadequate treatment are at higher risk for pregnancy-related relapses. Obstetric complications such as first-trimester miscarriage, preeclampsia and

preterm delivery complicate the course of pregnancy. However, it seems that the intensive treatment with corticosteroids and immunosuppressants can control most disease recurrence during pregnancy and the postpartum period. For severe cases, intravenous immunoglobulin, plasma exchange and immunoabsorption are employed, while the use of monoclonal antibodies during pregnancy is yet under debate. Since NMOSD cases are becoming more frequent worldwide, the management protocols are warranted to guide clinicians.

Keywords: neuromyelitis optica spectrum disorder, pregnancy, Aquaporin-4

Increase of the rates of premature birth and of caesarean section generated by COVID-19 in a tertiary maternity

Octavian Gabriel Olaru, Anca Daniela Stănescu, Ileana Conea, Denisa Oana Bălălau, Romina Marina Sima, Liana Pleș

Department of Obstetrics and Gynecology, "Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

Introduction. In the context of the COVID-19 pandemic, there are many controversies about the influence of this disease on pregnancy. More and more data are being collected on the risk of premature birth and the risks associated with birth. **Purpose of the study.** This study aimed to analyze the changes in the profile and management of obstetric cases with SARS-CoV-2 infection. **Materials and method.** This retrospective study, in a single tertiary reference center maternity, included all consecutive cases admitted to the Bucur Maternity Hospital, "Sf. Ioan" Emergency Clinical Hospital (Bucharest, Romania), infected with SARS-CoV-2, focusing on the cases that gave birth. Data regarding the medical condition and birth were extracted from medical records. **Results and discussion.** During one-year period, starting from March 2020, a total of 314 cases

infected with SARS-CoV-2 were hospitalized in the clinic. According to clinical classification criteria, 53% of cases were asymptomatic, 38% were mild, 5% were medium and 4% of cases were severe. There were 267 births. The rate of premature births calculated for cases with COVID-19 was 17%, a significant increase compared to the national and previous maternity rate (RR=2.82; 95% CI; 1.54 to 5.1, $p<0.001$). The rate of caesarean births in COVID-19 cases reached 88%, which represented a significant increase compared to previous data (RR=1.46, 95% CI; 1.32 to 1.62, $p<0.0001$). **Conclusions.** COVID-19 causes a significant increase in the rate of premature births. Multifactorial negative influence on the birth leads to an increase of the caesarean section rate.

Keywords: premature birth, caesarean, COVID-19, SARS-CoV-2

Cervical cancer screening in Romania: it is time to change

Gheorghe Peltecu, Anca Maria Panaitescu, Nicolae Gică

Department of Obstetrics and Gynecology, "Carol Davila" University of Medicine and Pharmacy, Bucharest; "Filantropia" Clinical Hospital of Obstetrics and Gynecology, Bucharest, Romania

Cervical cancer is one of the most frequent female cancers in the world, mainly in low-resource countries. It is preventable through vaccination before starting sexual activity. The progression from infection with high-risk (HR) strains HPV to invasive cancer takes time, from 10 to 13 years, or even longer. During this long natural history of precancer phase, early detection and local treatment of the screened population could stop the progression to invasive cancer and would make this infectious neoplasia a preventable disease. The highest incidence of invasive cervical cancer and the highest death rate due to this disease are encountered in low-resource countries

with no preventive care programs. The key of success is an organized screening program that could dramatically reduce the incidence and mortality from cervical cancer through early detection and local treatment of early precancer lesions. Romania has the highest rate of mortality due to cervical cancer in Europe and has no national screening program to prevent cervical cancer. Opportunist screening is substantial in urban areas, but it does not cover the rural zones. Romania urgently needs a nationwide screening program to detect precursor lesions of cervical cancer.

Keywords: cervical cancer, screening

Sentinel lymph node biopsy in malignant pathology of vulva

Gheorghe Peltecu, Anca Maria Panaitescu, Nicolae Gică

Department of Obstetrics and Gynecology, "Carol Davila" University of Medicine and Pharmacy, Bucharest; "Filantropia" Clinical Hospital of Obstetrics and Gynecology, Bucharest, Romania

The use of sentinel lymph node biopsy (SLNB) in the management of vulvar cancers (squamous and melanoma) represents the gold standard. It reduces the extent of the surgical procedure, offering prognostic information and reducing morbidity by overtreatment. When diagnosed, 30% of the vulvar cancers have metastases to the inguinal-femoral lymph nodes and 10% to the pelvic nodes. Surgery of vulvar cancers (including inguinal-femoral lymph nodes dissection) is associated with a high risk of infectious morbidity and wound dehiscence. SLNB offers prognostic information. The survival rate at five years after surgery is 64.9% in case of positive SLN on H&E stain and 92.1% for IHC. In case

of negative SLN, the recurrence rate is 3.6%. Current standards recommend complete lymph node dissection (CLND) in case of positive SLNB. All these recommendations are valid for squamous cancers, but not for vulvar melanoma. Recent studies have shown that, in case of positive SLNB, there are not benefits of CLND at three years. Thus, CLND is no more the gold standard in case of positive SLN in case of vulvar melanoma. A critical point in practicing SLNB is the learning curve of surgeons. Therefore, together the trasor(s) used and a clear protocol of each institution will offer the best results in treating the vulvar cancers.

Keywords: sentinel lymph node, vulvar cancer

Does birth improve the outcome of pregnancies with mild or severe SARS-CoV-2 infection? Our experience

Liana Pleș^{1,2}, Romina-Marina Sima^{1,2}, Tina-Ioana Bobei^{1,2}, Georgiana Răducanu², Anca-Daniela Stănescu^{1,2}

1. Department of Obstetrics and Gynecology, "Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

2. Department of Obstetrics and Gynecology, "Sf. Ioan" Hospital, "Bucur" Maternity, Bucharest, Romania

Introduction. SARS-CoV-2 pandemic had a worldwide enormous impact in healthcare and influenced every one's life. Pregnant women were also affected and the symptomatology varied from mild to severe forms. **Materials and method.** We present the experience of our clinic regarding the impact of delivery in pregnant patients with moderate and severe clinical forms of COVID-19. **Results.** On the 19th of March 2020, the "Bucur" Maternity became a center of care exclusively for pregnant women with SARS-CoV-2 infection. During one-year period, we had a number of 585 hospitalized pregnant women. Among them, 172 gave birth, with a 88.7% percentage of caesarean sections.

The indications of delivery by caesarean section were in 34.5% due to SARS-CoV-2 symptoms that imposed premature delivery. We report severe cases of acute respiratory distress symptoms and coagulopathies. The patients with moderate and severe respiratory failure imposed orotracheal intubations in 12 cases. In our clinic, it was observed that women with moderate or severe COVID-19 symptoms had a favorable outcome after delivery. **Conclusions.** In all cases that required caesarean section for severe respiratory distress, the maternal status improved, with supportive intensive care measures after delivery.

Keywords: COVID-19, delivery, respiratory distress

The impact of SARS-CoV-2 pandemic on obstetrics-gynecology residents' education

Liana Pleș^{1,2}, Romina-Marina Sima^{1,2}, Ioana Ciupitu², Anca-Daniela Stănescu^{1,2}

1. Department of Obstetrics and Gynecology, "Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

2. Department of Obstetrics and Gynecology, "Sf. Ioan" Hospital, "Bucur" Maternity, Bucharest, Romania

Introduction. The COVID-19 pandemic had a specific pattern in Romania regarding the hospitals' status. Thus, there were divided in COVID-19 hospitals and non-COVID-19 hospitals. The same distribution was made for maternities. This study aimed to identify the levels of education in obstetrics-gynecology residents from Romanian hospitals during the COVID-19 pandemic. **Materials and method.** We conducted a survey with a questionnaire completed by 85 obstetrics-gynecology residents aged between 25 and 32 years old, from COVID-19 and non-COVID-19 hospitals. **Results.** In both COVID-19 and non-COVID-19 hospitals, the level of education decreased during pandemic. The interviewed doctors agreed in a significant percentage of

65% that their practical part of preparation suffered, especially in COVID-19 hospitals, an answer approved by 97.5% of the responders. Residents considered that their decreased in preparation decreased due to the reorganization of work (67.6% for COVID-19 hospitals and 45.7% for non-COVID-19 hospitals). Overall, during this period, the responders answered that they received more theoretical information using online systems (78.5% in COVID-19 hospitals and 36.7% in non-COVID-19 hospitals). **Conclusions.** Among Romanian residents in obstetrics-gynecology, the COVID-19 pandemic was associated with significant impairment, especially in COVID-19 hospitals.

Keywords: residents, training, education, pandemic

The hysteroscopic management of abnormal uterine bleeding

Cristian Poalelungi^{1,2}, Iuliana Ceaușu^{1,2}, Nicoleta Alina Mareș¹, Maria Videnie¹

1. Department of Obstetrics and Gynecology, "Dr. Ion Cantacuzino" Clinical Hospital, Bucharest, Romania

2. "Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

Objective. Abnormal uterine bleeding is one of the main problems of diagnosis and evaluation in gynecological pathology. The current objective is the evaluation of the uterine cavity in the presence of endometrial polyps by transvaginal ultrasonography, followed by diagnostic and interventional hysteroscopy. **Methods.** The study included 41 cases of metrorrhagia hospitalized in the gynecology department of the "Dr. Ioan Cantacuzino" Clinical Hospital, Bucharest, during a period of a pandemic year. Our patients, women aged between 20 and 71 years old, were evaluated in the first stage by clinical examination and transvaginal ultrasound. In 25 cases, there were endometrial polyps between 1 and 3 cm in size, with persistent symptoms such as bleeding. All patients agreed to the hysteroscopic removal procedure. Exploratory hysteroscopy with endometrial biopsy was performed in the other cases without polyps. **Results.**

This study evaluated one of the most common pathologies that cause abnormal uterine bleeding and the importance of proper evaluation of the uterine cavity for the accuracy of diagnosis and therapeutic management. All women underwent biopsy and had a histological result, with: hysteroscopy followed by curettage, biopsy by hysteroscopy or hysteroscopy and resectoscopy. After procedure, there were moderate bleeding and less pain. **Conclusions.** The availability of advanced hysteroscopic techniques has changed the management of abnormal uterine bleeding. Along with careful clinical examination and ultrasonography, hysteroscopy seems to be the best option for a correct evaluation of the uterine cavity and for the accuracy of samples for the pathology exam of the endometrium.

Keywords: abnormal uterine bleeding, hysteroscopy, endometrial polyp, endometrial biopsy

Adolescent pregnancy: complications and birth rates

Daria Maria Pop¹, Renata Nicula¹, Cezarin Todea¹, Ciprian Porumb¹, Hognogi Radu Mocan¹, Ligia Blaga², Dan Mihui¹, Doru Diculescu¹

1. Second Department of Obstetrics and Gynecology, County Emergency Clinical Hospital Cluj-Napoca; "Iuliu Hațieganu" University of Medicine and Pharmacy, Cluj-Napoca, Romania

2. Department of Neonatology, "Iuliu Hațieganu" University of Medicine and Pharmacy, Cluj-Napoca, Romania

Introduction. Adolescent pregnancy has the potential to become a health issue in our country, especially since the birth rates in this category seem to be increasing. Teenagers who give birth are likely to have poor school performance and come from families with low socioeconomic status, a history of teen pregnancies, and low maternal education. Also, obstetrical literature describes low birth weight, perinatal death and preeclampsia associated with pregnancy and birth under the age of 18. The aim of this paper was to assess the pregnancy-associated risks as well as birth rates of adolescent pregnancies in Cluj-Na-

poca between 2019 and 2021. **Materials and method.** We reviewed data concerning pregnancy and births in adolescent mothers, aged 12 to 17 years old, who were admitted in our clinic from 2019 to 2021. Aside from age, we collected data regarding socioeconomic status, education, pregnancy-associated complications, way of delivery, parity and perinatal complications. **Conclusions.** Our findings strongly suggest that adolescent pregnancy is associated with environmental characteristics.

Keywords: adolescent pregnancy, adolescent birth, adolescent mother, teenage pregnancy

The usefulness of the SPIKES protocol in obstetrician-couple communication in case of detecting intrauterine fetal malformations

Dragoș Popescu^{1,2}, Radu Chicea^{1,2}, Iulia Lazăr²

1. "Lucian Blaga" University of Sibiu, Romania

2. County Emergency Clinical Hospital of Sibiu, Romania

Ultrasound technology has improved rapidly in the last years and its increased availability has made ultrasound detection of miscarriage, fetal abnormalities and of other pregnancy complications more common. The physicians' knowledge of how to give pregnancy-related bad news is becoming progressively important. A compassionate and informative method of breaking bad news, particular to the type of news, the anxiety level of the woman, the capacity to process and understand

the news for the woman, the physicians' comfort level and the resources that are available for emotional support are essential. Although studied in oncology, strategies for the communication of abnormal ultrasound information during pregnancy are scarce. We suggest a therapeutic approach to giving bad news during or after obstetrical ultrasound.

Keywords: prenatal ultrasound, fetal anomalies, obstetrician, bad news, protocol

Postoperative period management in patients after abdominal birth

Oleg Potacevski

Department of Obstetrics and Gynecology, "Nicolae Testemițanu" State University of Medicine and Pharmacy, Chișinău, Republic of Moldova

Introduction. The current problem of contemporary obstetric surgery is the increased risk of septic-purulent complications in the postoperative period, caused by the increased incidence of caesarean sections, especially those performed in emergency. **The aim of the study** is to identify conditions for sufficient healing of the uterus and to ensure the optimal conduct on the postoperative period. **Materials and method.** A retrospective study was performed on 88 patients, aged between 18 and 39 years old, who gave birth by urgent caesarean section, by transverse crescent hysterotomy in the lower segment, in the First Obstetrics Department of the Municipal Perinatalogical Center of Chișinău, Republic of Moldova, in 2020. **Results.** There were 36 repeated caesareans and 52 primary caesareans. The interventions took place under locoregional anesthesia in 78 cases (88.6%), and under endotracheal anesthesia in 10 cases (11.4%). The duration of operations varied from

30 minutes to 90 minutes. Hystero-graphy was performed with Vicryl® in 88.6% of cases and in 11.4% of cases with catgut. In the suturing technique, uninterrupted suturing prevailed in one layer (86.4% of cases). Preoperative antibiotic prophylaxis was performed in all patients (antibiotic therapy in 20.5% of cases). In all cases, early mobilization and enteral support with multimodal analgesia were practiced in the postoperative period. There were 10.2% of cases of complications in the postoperative period, in the form of postoperative hemorrhage, endometritis and suppuration of the postoperative wound. **Conclusions.** In order to reduce the incidence of postoperative complications in patients after abdominal births, it is necessary to reduce the duration of operations, standardize surgical technique and ensure an optimal postoperative management.

Keywords: caesarean section, postoperative complications

The influence of locoregional anesthesia on the evolution of the postoperative period

Oleg Potacevski

Department of Obstetrics and Gynecology, "Nicolae Testemițanu" State University of Medicine and Pharmacy, Chișinău, Republic of Moldova

Introduction. Locoregional anesthesia presents the method of choice (up to 90%) in contemporary surgical obstetrics, especially in connection with the increased incidence of caesarean operations. This ensures a favorable evolution of the postoperative period. **The aim of the study** was to analyze the influence of locoregional analgesia on the evolution of the postoperative period in patients after abdominal births. **Materials and method.** A retrospective study was performed on 88 patients, aged 18-39 years old, who gave birth by urgent caesarean section, by transverse crescent hysterotomy in the lower segment, in the First Obstetrics Department of the Municipal Perinatalogical Center of Chișinău, Republic of Moldova, in 2020. **Results.** There were 36 repeated caesareans and 52 primary caesareans. The interventions took place under locoregional anesthesia in 78 (88.6%)

cases (spinal in 64 cases and epidural in 14 cases), and under endotracheal anesthesia in 10 (11.4%) cases. The duration of operations varied from 30 minutes to 90 minutes. The presence in the ICU was between 12-36 hours, six patients needed hemotransfusion, and in all cases the early mobilization and the enteral support were performed with multimodal analgesia, in the postoperative period. Eighty-two patients presented moderate pain syndrome, while severe pain was present in only six patients. The duration of hospitalization varied between 25 and 72 hours. Three patients presented postanesthetic complications. **Conclusions.** Locoregional analgesia ensures a favorable evolution of the postoperative period in patients after caesarean section.

Keywords: caesarean section, locoregional anesthesia, postoperative period

Fetal shunt placement

Hadi Rahimian¹, Carmen Cristea¹, Oana Zaharia¹, Camelia Miron^{1,2}

1. "Regina Maria" Maternity, Bucharest, Romania

2. "Titu Maiorescu" University of Medicine, Bucharest, Romania

Fetal shunt placement involves the insertion of a shunt (tube) through the mother's abdomen into the uterus and into the fetus to drain fluid from a fluid filled space into the amniotic cavity. Vesicoamniotic shunt placement (used in posterior urethral valves syndrome) restores amniotic levels to normal by shunting fetal urine from the obstructed urinary system to the amniotic cavity, preventing lung hypoplasia and, thus, improving the neonatal survival. Reducing back pressure would also reduce nephron damage, improving postnatal renal function. Likewise, hydrocephalus due to severe ven-

triculomegaly causes permanent brain damage due to increased intracerebral pressure. Because of the obstruction of cerebrospinal fluid flow, CSF accumulates in the cerebral ventricles, causing increased intracerebral pressure, which leads to decreased blood flow, with neuronal damage. The intrauterine improvement of the increased intracerebral pressure by placing a ventriculoamniotic shunt can lead to a normal development of the brain, thereby preventing lifelong disability.

Keywords: fetal shunt placement, ventriculomegaly, posterior urethral valves syndrome

Twin-twin transfusion syndrome management

Hadi Rahimian¹, Carmen Cristea¹, Oana Zaharia¹, Camelia Miron^{1,2}

1. "Regina Maria" Maternity, Bucharest, Romania

2. "Titu Maiorescu" University of Medicine, Bucharest, Romania

STT occurs in monochorionic twin pregnancies due to the imbalance between the communicating vessels in the placenta, which leads to hypervolemia for one twin, with hypovolemia for the other. The management of twin-twin transfusion syndrome can be: expectant management, fetoscopic laser ablation of placental anastomoses, amnioreduction or selective fetal reduction. Fetoscopic laser ablation is a procedure by which a laser is inserted through the fetoscope and used to ablate the superficial blood vessels that cross the interamniotic membrane. Even though there are deep anastomoses, their afferent and efferent branches are superficial, so the coagulation of superficial vessels

should balance the interfetal transfusion. The complications of this technique are: premature birth, premature rupture of membranes, separation of membranes, rupture of interamniotic membranes, intraamniotic bleeding, fetal death, TAPS, persistent or recurrent TTTS, reversed TTTS. Quintero staging and correct clinical evaluation (significant respiratory distress, preterm contraction, cervix length) are essential for TTTS management. Untreated, 75% of stage I cases remain stable or regress, while perinatal loss for stages III-V is between 70% and 100%.

Keywords: twin-twin transfusion syndrome (TTTS), management

Indoor environmental quality assessment in a historical building – a maternity field-study in North-West Romania

Tania Rus¹, Gheorghe Cruciat², Georgiana Nemeti², Iulian Goidescu², Florin Vasile Domnița¹, Daniel Mureșan²

1. Faculty of Building Services Engineering, Technical University, Cluj-Napoca, Romania

2. First Obstetrics and Gynecology Clinic, Mother and Child Department, Emergency Clinic County Hospital, Cluj-Napoca; "Iuliu Hațieganu" University of Medicine and Pharmacy, Cluj-Napoca, Romania

Maintaining sustainability and proper indoor environmental quality (IEQ) in historical buildings is a challenging and ongoing issue across Europe. Historical buildings, compared to old buildings, are more complicated to be restored or retrofitted due to their historic value, cultural heritage and restoration policies. Another delicate aspect of the retrofit process is the achievement of a satisfactory level of IEQ, which plays a key role in the occupants' comfort, productivity, health and well-being. Indoor environmental quality comprises factors such as thermal comfort, air quality, acoustics and lighting quality. The aim of the research is to evaluate the IEQ of a tertiary level maternity located in a historical building dating from

1902 in North-West Romania. Field measurements of environmental factors were performed in obstetrics wards accommodating postpartum patients and their newborns during the summer season. Compliance with the current standards regulations of the recorded indoor environmental parameters was assessed. The results of the field measurements confirm the environmental thermal comfort factor which reached a satisfactory level in the study population. Lighting quality fulfils the standard recommendations. The mean recorded values of the acoustics and air quality exceeds the threshold limits.

Keywords: indoor environmental quality, thermal comfort, air quality, lighting, acoustic, hospital, maternity

Cervical remodeling in preterm birth. Timing, diagnosis and actual therapeutic opportunities

Manuela Russu^{1,2}, Adriana Ghelmen^{1,2}, Ruxandra Stănculescu², Șerban Nastasia^{1,2}

"Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

Department of Obstetrics and Gynecology, "Dr. Ion Cantacuzino" Clinical Hospital, Bucharest, Romania

Preterm birth (PTB) is a continuing burden for medical staff all over the world, including Romania. Premature cervical remodeling (CR) depiction and correction represent a prophylactic and/or a therapeutic option. Cervix uteri – either a metabolic active organ beginning from pregnancy or a piece of resistance/"gatekeeper" for normal evolution – is a prime source of early information on PTB risks, cervical softness and length changes being previous to preterm/term uterine contractions. The four successive stages – "softening, ripening, dilation, repair" – are at term, and "softening, ripening, dilation" can be the very early signals for late miscarriage/PTB, preterm rupture of membranes, thus their earlier depiction permits better prevention strategies. Longitudinal ultrasound and techniques of shear wave speed (SWS) assess the cervical stiffness from 8-13 weeks, showing a constant fractional reduction (about 4% per week) in SWS with the increasing of gestational age, and also a spatial gradient in SWS along the length of the cervix (softest at the distal end). When SWS technique is associated to serial dynamic transvaginal ultrasound cervical length (CL) measurements (at 16-23 weeks), they can accurately depict by multiple observations the subtle

cervical-segmental remodeling: sludge, CL, internal os dilation with funneling, percentage, form – T/Y/V/U, depth and width, protuberant aspect of inferior pole membranes, fetal parts/umbilical cord in the cervix/vagina, and allow the assessment of residual CL – cervical closed part (canal length), when membranes of inferior pole are ruptured, or with aspect of hourglass, when vaginal blood loss is active. CL ≤ 25 mm is associated to intraamniotic inflammation, and when CL ≤ 15 mm at 22-30 weeks, intraamniotic infection risks are higher. Vaginal progesterone, from early pregnancy (6-8 weeks) continued to 37 weeks, cerclage and pessary are actual therapeutic opportunities for a mechanical support, being discussed preterm births – history/actual, and elective/emergency – initial phase/urgency, when cervical incompetence. The cerclage is invasive, modern pessary is noninvasive, when cerclage cannot be recommended/contraindicated. Multicenter retrospective/prospective, randomized studies and meta-analyses are discussing them separately either progesterone or pessary adjuvant to cerclage.

Keywords: cervical remodeling, preterm birth, timing, progesterone, pessary, cerclage

High-risk HPV involvement in post-therapeutic CIN2+/CIN3+ recurrence

Manuela Russu^{1,2}, Cătălina Savu^{1,2}, Ruxandra Stănculescu², Anca Popa³, Șerban Nastasia^{1,2}

1. Department of Obstetrics and Gynecology, "Dr. Ion Cantacuzino" Clinical Hospital, Bucharest, Romania

2. "Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

3. Medsana Medical Centre, Bucharest, Romania

Contemporary medicine sustains HPV high risk (HPVhr) involvement in the development of extragenital (women and men) and lower anogenital cancer precursors, and their progression to invasive cancers. Even the studies are demonstrating the human clearance capacity; HPVhr is persistent after the current therapies for cervical intraepithelial neoplasia (CIN). The last systematic review (2018) presents two types of cervical HPV persistent infections: one that persists after the treatment for a previous lesion, and a new infection, post-therapy acquired. One discusses women's age, HPV status and genotype diagnosis, the procedure for treatment (conization, LLETZ versus cryotherapy, laser therapy, interferon alpha, therapeutic vaccination, and photodynamic therapy), searching data for vaccination efficacy, for the best clinical practice. Globally, it is described a gradual clearance after CIN treatment. In 2014, the median HPV persistence tended to decrease with increasing follow-up time, declining from 27% at three months after the treatment to 21% at six months, 15% at 12 months, and 10% at 24 months, compared to an old report, from 1994, of 100% persistence at 6 and

at 12 months. There are many arguments for HPVhr persistence and for the future risk of invasive cervical carcinoma in women treated for CIN2/3: many years since the intervention, women's age – higher risk in women above 50 years old, ethnicity, genetic variations in E6, E7 HPVhr oncoproteins, patients' viral quantification, the interactions between viral DNA and host DNA, as host DNA machinery permits/does not permit viral replication, infection's cofactors (other STDs – bacterial, including *Mycobacterium tuberculosis*, viral, including HIV and HVS; and general host polymicrobial infections – oral, digestive, respiratory, which reduce the patient immune competence), toxic factors (smoking), epigenetic changes (DNA methylation, progression from CIN2/3 to invasive carcinoma by epigenetic switch or by molecular switch), the long use of combined oral contraceptives (16 alpha-hydroxylation of estrogens from COC increases the transcription of HPVhr when HPVhr positive).

Keywords: HPVhr persistence, CIN2+/CIN3+, recurrence, viral quantification, DNA methylation, molecular switch

The place of colposcopy in modern gynecology. Indications, logistics and procedures

Ioan Sas^{1,2}, Larisa Tomescu^{1,2}, Maia Todorov¹

1. Department of Obstetrics and Gynecology, "Victor Babeș" University of Medicine and Pharmacy, Timișoara, Romania

2. Department of Obstetrics and Gynecology, County Clinical Emergency Hospital Timișoara, Romania

Introduction. World Health Organization confirms an incidence of 570,000 new cases of cervical cancer diagnosed in 2018 and a total of 270,000 deaths from this cause in 2015. These data rank cervical cancer as the fourth leading cause of death among the female population, representing 6.6% of all cancers in women. Approximately 90% of these deaths occurred in developing countries, and 95-98% of these lesions are caused by HPV (human papillomavirus) infection, with cases of increased oncological potential. Due to the high incidence, it is important to discuss and summarize the diagnostic strategy for preinvasive cervical lesions. **Methodology.** The aim of this paper is to demonstrate that colposcopy, introduced in 1925, and which is developing before the great technological advances in modern gynecology, continues to be a valid technique without essential innovations of the original method described at the beginning of the last century. A starting point for this effort was to define the role of colposcopy as a test used in the pre-

vention of cervical cancer. As with any screening, triage or diagnostic procedure, the risks and benefits of a test must be assessed and weighed. Another major component of this effort was the revision and standardization of terminology for colposcopy practice. The aim was to simplify and clarify the reporting of colposcopy findings to improve absorption by colposcopists practicing in diverse work environments. **Results and conclusions.** Colposcopy is an important step in the initial evaluation of the abnormal results of the cervical cancer screening test, and colposcopy should be considered a routine technique in daily practice. Digital colposcopy aims to improve cervical cancer screening and the diagnostic process, shortening the period between the first medical interaction and the implementation of treatment, helping diagnose preneoplastic and neoplastic cervical lesions and reducing mortality.

Keywords: digital colposcopy, cervical cancer screening, guideline recommendations

Second victims in obstetrics – a literature review

Ioan Sas^{1,2}, Nicoleta Nicolae^{1,2}, Denis Șerban^{1,2}, Larisa Tomescu^{1,2}

1. Department of Obstetrics and Gynecology, "Victor Babeș" University of Medicine and Pharmacy, Timișoara, Romania

2. Department of Obstetrics and Gynecology, County Clinical Emergency Hospital Timișoara, Romania

Medical errors are a serious public health problem. Every day, healthcare professionals practice their occupation within excessively complex healthcare situations and sudden patient outcomes. This paper aims to explore the knowledge about the phenomenon of the second victim in the obstetrics specialty. In 2000, the "second victim" was defined for the first time by Albert Wu in his description of the impact of errors on healthcare professionals by both personally and professionally, being a medical emergency equivalent to post-traumatic stress disorder. The term acknowledges the "first victim" to be the patients and, sometimes, the term "third victim" refers to the healthcare organization. The terminology has been criticized by patient advocacy communities for incorporating the word "victim", mainly as they use this term to refer only

to the patient. Medical errors, non-error patient safety events, near misses are common and can impact the well-being of the provider. This can lead to secondary traumatic stress such as emotional distress, sleep difficulties, anxiety, distress, PTSD, guilt/shame, fear, suicidality, and to negative effect on work performance such as absenteeism, reduced confidence and/or potential secondary medical errors. The recent recognition and discussion on the phenomenon of the second victim have become a popular research subject. This review intended to study not only the phenomenon of second victim in general medical care, but to also to concentrate on obstetrical care where the expectation of perfection may be argued to be greater.

Keywords: health personal, second victim, obstetric care

Current clinical and therapeutic challenges in HELLP syndrome

Romina-Marina Sima^{1,2}, Gabriel Gorecki^{2,3}, Andreea Marcu², Liana Pleș^{1,2}

1. Department of Obstetrics and Gynecology, "Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

2. Department of Obstetrics and Gynecology, "Sf. Ioan" Clinical Hospital, "Bucur" Maternity, Bucharest, Romania

3. Medicine Doctoral School, "Titu Maiorescu" University, Bucharest, Romania

Introduction. HELLP syndrome is described as hemolysis, elevated liver enzymes and low platelets, and it has been historically classified as a complication or progression of severe preeclampsia. **Materials and method.** We reviewed the recent reports from international data bases regarding the actual therapies and recommendations about HELLP syndrome. **Results.** The medical publications from the last years report that there is no direct relation between severe arterial hypertension and HELLP syndrome. This syndrome is a unique entity that may occur in normotensive patients. The genetic analysis of the inheritability for preeclampsia and/or HELLP syndrome in pregnancy has also been explored. The results present both genetic and immunological factors that have a determinant role in pathogenesis. The differential diagnoses for HELLP include

acute fatty liver of pregnancy, purpura, antiphospholipid syndrome, thrombotic thrombocytopenic and hemolytic uremic syndrome. There is not any current treatment for HELLP, but the mainstay of therapeutic approach involves maternal stabilization and timely delivery. Different treatment strategies have been attempted to help decrease the morbidity and mortality, such as the maternal use of corticosteroids. However, the studies present controversies regarding the maternal use of corticosteroids, plasma exchange, and low molecular weight heparin for the treatment of HELLP. **Conclusions.** The pathology of HELLP syndrome is far to be completely described and the treatment represents a challenge for physicians. The discussions about the topic are always with benefits in order to improve the patients' outcome.

Keywords: HELLP, arterial hypertension, treatment

The pathological characteristics of myometrium in uterine atonia – case report

Romina-Marina Sima^{1,2}, Cristina-Diana Pomană^{1,2}, Liana Pleș^{1,2}, Cătălina Alexandrescu³

1. Department of Obstetrics and Gynecology, "Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

2. Department of Obstetrics and Gynecology, "Sf. Ioan" Clinical Hospital, "Bucur" Maternity, Bucharest, Romania

3. Medlife Memorial Hospital, Bucharest

Introduction. Postpartum hemorrhage represents the leading cause of maternal mortality worldwide. Uterine atonia is one of the main causes and represents a therapeutic challenge for obstetricians. **Case report.** We report the case of a 34-year-old patient who was admitted in our clinic at 41 weeks of gestation for fetal monitoring. From her medical history, we mention an abortion with twin pregnancy at 18 weeks of gestation. The actual pregnancy was monitored according to our national recommendations and had no pathological event. The patient had no uterine contraction during pregnancy, not even at term. Due to unfavorable Bishop score at 41 weeks of gestations, the caesarean section was performed and a 4100 g female fetus with 10 APGAR score was born. The caesarean section developed with no significant hemorrhagic event (450

ml blood loss). The uterus retracted with 100 mcg of carbetocin. At three hours after birth, a vaginal severe hemorrhage occurred, with 2000 ml blood loss and the postpartum uterine atonia was diagnosed. It imposed explorative laparotomy. The medical and surgical techniques of uterine preservation (B-Lynch sutures) failed and supracervical hysterectomy was performed. The maternal evolution was favorable, with discharge four days after delivery. The pathological exam of the uterus revealed important uterine edema and atypical myometrial cells. **Conclusions.** Uterine atonia is a condition that is difficult to be anticipated, but it may be considered in patient with no uterine contractility near term, fetal macrosomia and significant myometrial architecture anomalies.

Keywords: uterine atonia, hemorrhage, edema

Management challenges of psoriasis in pregnancy

Anca Angela Simionescu

Department of Obstetrics and Gynecology, "Carol Davila" University of Medicine and Pharmacy, Bucharest; "Filantropia" Clinical Hospital of Obstetrics and Gynecology, Bucharest, Romania

Psoriasis is an immune-mediated inflammatory disease with genetic predisposition, manifested as skin and joints features. There are many challenges in the management of psoriasis in pregnancy due to the interaction with fetomaternal immune system barrier, psoriasis comorbidities and limited options, and conflicting and limited data regarding drugs used in pregnancy. The immunology of pregnancy is unique and characterized by maternal immune response changes from the inflammatory Th1 cytokine pattern to the Th2 pattern. These cytokines are critical for the adaptive immune responses during pregnancy. The dysfunction of T-cell subsets of Th1 cells is responsible for psoriasis lesions and the resulting aberrant release of the corresponding cytokines, including IFN- γ , tumor necrosis factor (TNF)- α , IL-23 and IL-17 family members. The immunology of pregnancy is related to CD4⁺ T cell cytokines, and T cell responses in autoimmune disease are influenced by pregnancy. In more than 40% of reported

cases, psoriatic skin and joints lesions improved during pregnancy, in 20% of cases the evolution worsened, and 21% to 56% remained stable. Psoriasis is associated with comorbidities such as diabetes mellitus, metabolic syndrome, depression and an increased risk of alcohol and tobacco intake; this can negatively influence pregnancy evolution and outcomes. The classical treatment regimens for psoriasis in pregnancy include low-doses of topical and systemic therapies and, recently, human monoclonal antibodies for moderate to severe forms. Secukinumab and ustekinumab are becoming interesting options to the classical regimen, despite the high costs and the limited experience during pregnancy. The first-line therapy used for women of childbearing age remains certolizumab pegol, a PEGylated, Fc-free anti-TNF agent that does not cross the placenta. In conclusion, moderate to severe psoriasis forms during pregnancy remain a very high risk to manage.

Keywords: psoriasis, pregnancy, immune response

Epidemiology of clavicle fracture at birth in Romania

Anca Angela Simionescu¹, Vlad Dima²

1. Department of Obstetrics and Gynecology, "Carol Davila" University of Medicine and Pharmacy, Bucharest; "Filantropia" Clinical Hospital of Obstetrics and Gynecology, Bucharest, Romania

2. Department of Neonatology, "Filantropia" Clinical Hospital of Obstetrics and Gynecology, Bucharest, Romania

A clavicle fracture is the most common unpredictable bone trauma at birth; most cases occur following uncomplicated vaginal deliveries. The reported incidence is 0.2-2.9% of births, mostly diagnosed at a distance from childbirth. This trauma at birth can be seen by parents as a medical error of birth care, especially when diagnosed at a distance from hospitalization. The purpose of this poster is to present Romanian statistics in the last two years from the perspective of the obstetrician and neonatologist. In 2018, 358 cases of clavicle fracture were reported for 163,490 births and, in 2019, 308 cases of clavicle fractures for 1,585,20 births, with an incidence of 0.19-0.21% of births, well below the one reported by other countries. A clavicle fracture in a newborn can be difficult to diagnose because it is often asymptomatic and can also be confused with other common diagnoses, such as brachial plexus palsy, congenital pseudoarthrosis and congenital muscular torticollis.

The common causes are shoulder dystocia or breech presentation deliveries, when the arms of the baby are extended. Evidence from retrospective studies demonstrates that excessive weight gain by mother and labor induction may increase the risk of clavicular fracture with brachial plexus palsy lesions. A greenwood fracture or an incomplete fracture may be asymptomatic at birth. The first clinical sign may be callus formation 7 to 10 days after birth. The clinical sign often identified shortly after birth is edema in the supraclavicular fossa, with underlying crackles in general. The palpation of the clavicle should always be repeated on the discharge examination, as some fractures may be more obvious on the second or the third day of life.

Acknowledgements: We want to acknowledge Lucia Dragomir and Ecaterina Matei from the National Institute of Public Health for these data.

Keywords: clavicle fracture, vaginal delivery, trauma

Placental pathologic changes in gestational diabetes mellitus and implications for fetal outcome

Adelina Staicu, Héloïse Pelczar, Camelia Albu, Ioana Cristina Rotar, Andreia Preda, Roxana Constantin, Iulian Goidescu, Dan Boitor-Borza, Mihai Surcel, Gheorghe Cruciat, Daniel Mureșan

First Obstetrics and Gynecology Clinic, Emergency County Clinical Hospital Cluj-Napoca; "Iuliu Hațieganu" University of Medicine and Pharmacy, Cluj-Napoca, Romania

The placenta is a transitory organ which supports intrauterine life with nutritional, endocrine and immunologic functions, whose adequate functioning is necessary for the development of the *fetus in vitro*, but it also impacts the biochemical mechanisms in the long run. Pathologies developed during pregnancy, such as gestational diabetes mellitus, influence the function and structural aspect of the placenta which in turn presents a number of alterations. The aim of our research was to compare the histopathological findings of placenta and umbilical cord between patients diagnosed with gestational diabetes mellitus (GDM) and physiological pregnancies (non-GDM). **Materials and method.** In this retrospective study, 110 patients were included (29 GDM and 81 non-GDM). GDM was confirmed at 24-28 weeks of gestation, using oral glucose tolerance test. We analyzed the demographic characteristics of the patients and the histologic and gross morphology findings of placentas. **Results.** The mean age at birth of GDM group was 31.38 years old, compared to 29.7 years old for non-GDM ($p=0.06$), and the mean gestational age at

birth was 37.41 weeks of gestation (WG) for GDM, compared to 38.5 WG in the non-GDM group ($p=0.006983$). The histological evaluation revealed significant larger placentas in the GDM group, with more inter- and perivillous fibrinoid ($p<0.0001$), intervillous thrombus ($p=0.0012$; 6.58-2060.72; OR=116.42), increase villous maturation ($p<0.0001$; 32.01-1048.92), chorangiosis ($p<0.0069$; 2.988-988.021), villous ischemia ($p<0.0001$; 12.0136-806.971), villous hypoplasia ($p<0.0001$; 32.01-10848.917), and hyperspiral cord ($p<0.00001$). Regarding calcifications, placental hematomas, placental infarction and chorioamnionitis, we found no relationship between their presence and gestational diabetes. **Conclusions.** This study highlights a series of macroscopic and microscopic placental changes in patients diagnosed with GDM: increased inter- and perivillous fibrinoid, important intervillous thrombus, accelerated villous maturation, chorangiosis, villous ischemia and hypoplasia and hyperspiral cord.

Keywords: placenta, gestational diabetes mellitus, fetal outcome

Update of the epidemiology and ultrasound diagnosis criteria able to predict before delivery *placenta accreta* spectrum disorders

Ruxandra Stănculescu¹, Manuela Russu^{1,2}, Elvira Brătîlă^{1,3}, Vasilica Bausic¹, Oana Toader^{1,5}, Ciprian Coroleucă^{1,3}, Alexandra Bausic³, Patru Papuc⁴, Ramona Dragomir⁵

1. Department of Obstetrics and Gynecology, "Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

2. "Dr. Ion Cantacuzino" Clinical Hospital, Bucharest, Romania

3. "Prof. Dr. Panait Sirbu" Clinical Hospital of Obstetrics and Gynecology, Bucharest, Romania

4. "Sf. Pantelimon" Emergency Clinical Hospital, Bucharest, Romania

5. "Polizu" Hospital, "Alessandrescu-Rusescu" National Institute for Mother and Child Health, Bucharest, Romania

Introduction. The increasing incidence of abnormal insertion of placenta justifies the need to highlight which are the frequent factors involved in this pathology and to improve the methods able to sustain a prenatal diagnosis of abnormal placenta spectrum disorders, so useful to avoid a catastrophic postpartum hemorrhage. **Material and method.** The study was conceived by analysis of articles and clinical guidelines published online concerning the *placenta accreta* spectrum disorders (PAS) in the last five years. The study allows to clarify the terms that define PAS, the factors that frequently generate this pathology, and the ultrasonographic criteria able to support PAS diagnosis. **Results.** The most proper terminology remains the one launched by Luke (1966), because it includes all three significant criteria, such as the depth of the myometrial invasion of the villi, the lateral extension of the *placenta accreta*, and the variable degrees of myometrial invasion in the same placenta. Among the factors most frequently involved, there are repeated caesarean sections, uterine curettages and manual extraction of the placenta. The

features of ultrasonographic criteria were the loss of hypoechogenic space, placental abnormal lacunae, loss or disruption of bladder wall continuity, thinning of the thickness of the myometrium, bulging of the placenta, focal placental exophytic, color Doppler changes such as uterovesical hypervascularization, subplacental hypervascularization, "bridging vessels" and Doppler intramiometrial vessels with high velocimetry oriented to the placental lacuna. The three-dimensional (3D) power Doppler sonography shows placental hypervascularization, vessels with a twisted path and different caliber. MRI is recommended to validate if a posterior placenta is *accreta* too. **Conclusions.** Women with risk factors for PAS should be screened by US at 11-14 weeks of pregnancy. The US criteria for antenatal diagnosis of SAP must be checked in the second trimester of pregnancy to avoid a catastrophic postpartum hemorrhage associated with disseminated intravascular coagulation syndrome.

Keywords: *placenta accreta* spectrum disorders, ultrasound, antenatal diagnosis, postpartum hemorrhage

The impact of SARS-CoV-2 infection on fluid-coagulant status of pregnant women

Anca-Daniela Stănescu, Romina-Marina Sima, Liana Pleș

Department of Obstetrics and Gynecology, "Carol Davila" University of Medicine and Pharmacy, Bucharest; Department of Obstetrics and Gynecology, "Sf. Ioan" Clinical Hospital, "Bucur" Maternity, Bucharest, Romania

Introduction. Physiological changes during pregnancy have an essential role on the cardiovascular function, the respiratory system and coagulation. There has been a rapid increase in publications about COVID-19 and pregnancy. The purpose of our research was to highlight the impact of SARS-CoV-2 infection on pregnancy and on the fluid-coagulant status. **Materials and method.** We searched the international databases using keywords such as "pregnancy", "COVID-19" and "coagulopathy". There were extracted only the publications about the implications of SARS-CoV-2 in pregnancy-associated coagulopathies. **Results.** The initial research revealed 3120 reports about pregnancy and COVID-19. Among them, the number of articles about the impact of SARS-CoV-2 in pregnancy fluid-coagulant status is reduced. There are

one meta-analysis and five case reports. The other studies about pregnancy and SARS-CoV-2 infection also present some patterns about this viral infection on pregnancy hemostasis. It was identified that coagulopathy or thrombotic complications may develop. Hypofibrinogenemia (compared to normal pregnancy levels) was identified in three cases. There is few evidence in the literature regarding platelet count in COVID-19 affected pregnancies. **Conclusions.** There is a lack of studies regarding the impact of SARS-CoV-2 infection in pregnancy fluid-coagulant status. The literature is based on case reports, but the clinical features presented are substantial. Further studies are required regarding the subject.

Keywords: SARS-CoV-2, fluid-coagulant status, pregnancy

Short-term disease recurrence after loop electrosurgical excision procedure (LEEP) – retrospective study

Irina Stoian^{1,2}, Ciprian Ilea^{1,2}, Demetra Socolov^{1,2}

1. "Cuza Vodă" Clinical Hospital of Obstetrics and Gynecology, Iași, Romania

2. "Grigore T. Popa" University of Medicine and Pharmacy, Iași, Romania

Introduction. Cervical cancer screening aims to identify cervical dysplastic lesions for the subsequent excision to prevent invasive disease. The surveillance of women with cervical dysplasia after treatment is performed by colposcopy or cervical cytology correlated with human papillomavirus (HPV) testing. The treatment of preinvasive disease is performed using loop electrosurgical excision (LEEP). The patients are typically followed-up for 12 months after the excision of the lesion. This study aims to quantify the prevalence of different types of cervical injuries in the region of Moldova, as well as the risk of recurrence following surgical excision. **Materials and method.** We retrospectively analyzed 308 patients (mean age: 36 years old) with pathologic Pap smear who were treated with LEEP, between 2010 and 2020. Post-LEEP follow-up was performed by Pap smear and colposcopy. Descriptive and analytic statistics were performed. **Results.** Of the 308 patients, 88.63% met the inclusion criteria. After Pap smear, high-risk lesions were found in 34.34% of the patients and low-risk lesions were identified in 50.97%

of them. HPV was performed in only 36% of cases. The anatomopathological results highlighted the following types of lesions: condyloma (1.62%), CIN 1 (24.68%), CIN 2 (16.23%), CIN 3 (10.39%), CIS (3.25%), microinvasive carcinoma (0.97%), invasive carcinoma (0.65%), and benign lesions (25.65%). During follow-up, 24 patients were lost from the records, and the control was performed on 100 patients. There were 41 cures, 52 recurrences and seven residual injuries. The percentage of discrepancies between the results obtained by screening and those resulting from the pathological examination was 20.77%. **Conclusions.** Over 50% of the lesions identified at screening were low risk. There is an increased percentage of recurrences following LEEP, especially in those with HPV-positive status. A significant predictive factor for persistent/recurrent disease was represented by margin and HPV status. A subsequent control and even a new excision are necessary to prevent the invasive disease.

Keywords: cervical dysplasia, follow-up, human papillomavirus, loop electrosurgical excision procedure

Magnesium sulphate and fetal neuroprotection – when, how, why and why not?

Viorela-Elena Suci¹, Răzvan Ciortea^{1,2}, Carmen Elena Bucuri^{1,2,3}, Andrei Măluțan^{1,2}, Radu Mocan-Hognogi^{1,2}, Maria Rada^{1,2}, Marina Ducea^{1,2}, Adelina Clim¹, Dan Mihu^{1,2}

1. "Dominic Stanca" Obstetrics-Gynecology Clinic, Emergency County Clinical Hospital, Cluj-Napoca, Romania

2. "Iuliu Hațieganu" University of Medicine and Pharmacy, Cluj-Napoca, Romania

3. "Constantin Papilian" Military Emergency Clinical Hospital, Cluj-Napoca, Romania

Preterm delivery is strongly associated with neurological impairments, such as cerebral palsy (CP), motor dysfunction, blindness, developmental delay and intellectual impairment. Antenatal administration of magnesium sulfate ($MgSO_4$) has become a valuable approach of the neuroprotective strategy for preterm births. $MgSO_4$ can prevent excitotoxicity via N-methyl-D-aspartic acid (NMDA) receptor antagonistic action and has an anti-inflammatory effect. There is still no international consensus on the dosage, time of administration and the need to repeat $MgSO_4$ treatment. However, most studies report benefits for $\leq 31+6$ weeks of gestation (GW) and imminent preterm birth, defined as active labor with ≥ 4 cm cervical dilation with either failure or contraindication to tocolysis, ≥ 4 cm dilatation with documented progressive change in cervical dilation, preterm pre-labor rupture of membranes with active labor, and for planned delivery for fetal or maternal indications. $MgSO_4$ is recommended

in a loading dose of 4 g intravenously (i.v.) for 30 minutes, followed by a 1 g/hour i.v. maintenance infusion for maximum 24 hours. It has been shown a significantly reduced rate of moderate and severe CP for the premature receiving $MgSO_4$, but also a lower rate of motor dysfunction and a lower systemic vascular resistance and higher myocardial function in preterm infants born before 29 WG. On the other hand, several side effects have been reported and should be closely monitored. Maternal side effects include hypotension, tachycardia, respiratory depression, drowsiness, headache and muscle weakness. One trial reported a higher incidence of spontaneous intestinal perforation among extremely low birth weight infants. In conclusion, antenatal $MgSO_4$ administration is a first step in preventing neurological damage, but further studies are mandatory to standardize its use.

Keywords: magnesium sulphate, neuroprotection, cerebral palsy

Therapeutic solutions for infertile “poor responder” patients

Mihai Surcel

Emergency County Clinical Hospital Cluj-Napoca; “Iuliu Hațieganu” University of Medicine and Pharmacy, Cluj-Napoca, Romania

Infertile “poor responders” represent a consistent group of patients addressing *in vitro* fertilization (IVF) programs. Despite tremendous efforts including the use of various ovarian stimulation and hypophysis control regimens, incremental gonadotropins doses or new drug molecules, IVF results are frustrating for both doctors and patients. This is mainly due to the plethora of factors that may be responsible for this type of infertility entity,

starting with intrinsic oocyte anomalies, problems concerning early phases of folliculogenesis or gonadotropin receptor dysfunctions. Attempts for a tailored approach of the different subtypes of poor responder patients resulted in the elaboration of the Poseidon criteria. New patient stratification and management strategies were developed, but the results are still under evaluation.

Keywords: poor responders, IVF, oocytes, gonadotropins

Ultrasound aspects of uterine scars after caesarean section in patients with retroverted uterus versus patients with anteverted uterus

Lucian Șerbănescu¹, Paris Ionescu¹, Diana Badiu¹, Dan Costea²

1. Department of Obstetrics and Gynecology, Faculty of Medicine, “Ovidius” University of Constanța; “Sf. Apostol Andrei” Emergency County Clinical Hospital, Constanța, Romania

2. Department of Surgery, Faculty of Medicine, “Ovidius” University of Constanța; “Sf. Apostol Andrei” Emergency County Clinical Hospital, Constanța, Romania

The study was performed at the Santerra Medical Center, Constanța, on two groups of patients. The first group included 100 pregnant patients with ultrasound-determined anteverted uterus before pregnancy, and the second group included 50 pregnant patients with ultrasound-determined retroverted uterus, before pregnancy. All patients studied were primiparous, who gave birth by caesarean section performed between 38 and 40 weeks of pregnancy (full term). Ultrasound aspects of uterine

scars at six weeks after caesarean section were analyzed (i.e., scar dimensions, presence of “L sign”). The incidence of larger scars (i.e., over 1.2 cm) and the presence of the L sign were significantly increased in patients from the second group. Uterine retroversion, through the mechanical effect it exerts on the post-caesarean hystero-graphy sheet, can cause a slower and sometimes even vicious scarring.

Keywords: caesarean section, primiparous, transvaginal ultrasound

Incidence of premature birth threat in patients with fibromatous uterus

Lucian Șerbănescu¹, Paris Ionescu¹, Diana Badiu¹, Dan Costea²

1. Department of Obstetrics and Gynecology, Faculty of Medicine, “Ovidius” University of Constanța; “Sf. Apostol Andrei” Emergency County Clinical Hospital, Constanța, Romania

2. Department of Surgery, Faculty of Medicine, “Ovidius” University of Constanța; “Sf. Apostol Andrei” Emergency County Clinical Hospital, Constanța, Romania

The study was performed in Clinic II of Obstetrics-Gynecology of the “Sf. Apostol Andrei” Emergency County Clinical Hospital, Constanța, on a group of 120 patients with uterine fibroids associated with pregnancy. Patients diagnosed with uterine fibroids after 24 weeks of gestational age were included in the study. The dimensions of the fibromatous nodules varied between 2 and 5 cm. All patients received progesterone therapy with intravaginal administration, 400 mg/day, in two doses, at 12 hours, until the gestational age of 36 weeks. The incidence of premature births in the studied group ver-

sus the general population was analyzed. No significant variations were found between the incidence within the studied group and the incidence in the general population. More than half of the patients in the study group complained of unexplained pelvic pain, uncorrelated with painful uterine contractions. It appears that progesterone given to patients with uterine fibroids associated with pregnancy helps to significantly reduce the incidence of preterm birth.

Keywords: premature birth, uterine fibroids, pregnancy

The impact of autoimmune, inflammatory and metabolic disorders on placental inflammation and perinatal outcomes

Atakan Tanacan

Perinatology Clinic of Maternity Hospital, Turkish Ministry of Health, City Hospital, Ankara, Turkey

Favorable pregnancy outcomes depend on optimal regulation of the maternal-fetal interface (MFI). Placental inflammation (PI) and impaired fetal perfusion are most likely attributable to the destruction of cellular components of the intervillous space (superficial decidual epithelium, endothelium of spiral veins, syncytiotrophoblasts of chorionic villi, and endovascular trophoblasts covering the tips of spiral arteries, among others) by various molecules/agents. The pathophysiological mechanisms that cause MFI injury result in PI, which contributes to obstetric complications and to poor pregnancy outcomes. Injury to cellular components of the MFI due to immune-mediated complexes and/or toxic metabolic components leads to PI. Cellular debris from the destruction of syncytiotrophoblasts and endovascular trophoblasts contains paternal antigens that may stimulate the maternal innate and humoral immune systems. Such processes alter the

immunologic balance of MFI and cause PI. Infectious, toxic, metabolic and immunological inflammation may be the underlying factors behind PI. Methylenetetrahydrofolate reductase polymorphisms, autoimmune/inflammatory diseases, metabolic disorders and hereditary thrombophilia are the most common risk factors behind the development of PI. There are recent publications in the literature indicating the impact of the etiology-based approach for preventing PI-associated obstetric complications and poor pregnancy outcomes. Lifestyle modifications, specific anti-inflammatory medications and strict antenatal follow-up protocols seem to reduce PI. In conclusion, the appropriate management of maternal risk factors and preventing PI should be the main targets of maternal-fetal medicine specialists.

Keywords: maternal-fetal interface, obstetric complications, placental inflammation

Conotruncal congenital heart diseases: prenatal diagnosis and management

Atakan Tanacan

Perinatology Clinic of Maternity Hospital, Turkish Ministry of Health, City Hospital, Ankara

Conotruncal malformations constitute an important part of congenital heart diseases. They include anatomic anomalies of both the infundibulum (conus arteriosus) and the great arteries (truncus arteriosus). Tetralogy of Fallot, *truncus arteriosus*, transposition of the great arteries and double outlet right ventricle are the most common conotruncal malformations in daily clinical practice. The accurate diagnosis and the appropriate antenatal management of these anomalies are vital to achieving favorable outcomes. With developing imaging technologies and increasing knowledge in the field of maternal-fetal medicine, it may be possible to diagnose these anomalies at relatively earlier weeks of gestation. If they are diagnosed prenatally, patient counseling by a multidisciplinary

team can be possible and necessary procedures can be implemented. The investigation of chromosomal/structural abnormalities, planning the delivery in a tertiary healthcare facility, organizing the neonatology team, arranging necessary cardiovascular interventions promptly and offering the option for termination of pregnancy in severe cases can be organized by maternal-fetal medicine centers. Moreover, the underdiagnosis of these anomalies may lead to medicolegal problems. For this reason, physicians dealing with obstetrics should be meticulous about the evaluation of the fetal heart and should carry out the necessary consultations at the slightest suspicion.

Keywords: congenital heart diseases, conotruncal malformations, prenatal diagnosis

Management of gastroesophageal reflux during pregnancy and lactation

Vlad Tica^{1,2,3,4}, Roxana Penciu^{1,2,3}, Liliana Steriu^{1,2,3}, Silvia Izvoranu^{1,2,3}, Corina Nour^{1,2,3}, Mihai Banariu^{1,2}, Diana Mocanu^{1,2,3}, Diana Badiu^{1,2,3}, Irina Tica^{1,3}

1. Faculty of Medicine, "Ovidius" University of Constanța, Romania

2. Institute of Doctoral Studies, Doctoral School of Medicine, "Ovidius" University of Constanța Romania

3. "Sf. Apostol Andrei" County Emergency Clinical Hospital, Constanța, Romania

4. Academy of Romanian Scientists

The clinical picture of gastroesophageal reflux disease (GERD) is relatively typical, with regurgitation and/or sensation of retrosternal burning, continuous or discontinuous. GERD is frequently met in pregnancy. In Romania (without national statistics), the prevalence estimate is 30-50%; globally, between two and eight in 10 women have transient GERD-specific symptoms. A small percentage of pregnant patients with GERD are treated appropriately, a context in which their quality of life is negatively affected. Some patients receive acid suppression therapy with proton pump inhibitors (PPIs). Despite treatment, some patients continue to experience reflux symptoms. The presentation reviews therapeutic options and current developments in GERD from pregnancy. Thus, international guidelines rec-

ommend changing your lifestyle as a first option. In case of persistent symptoms, the use of non-systemic drugs, such as antacids, is recommended as a first resort. Other therapeutic options, in the absence of a favorable result, are H2 receptor antagonists and, subsequently, PPIs – with significant adverse reactions for the long-term administration. Taking into account the degree of transplacental transport of the active substance and the teratogenic risk to the fetus, depending on the patient's needs, drug combinations are also considered in order to obtain the fastest effect, the longest duration of action and minimal effects on pregnancy and the fetus.

Keywords: gastroesophageal reflux, pregnancy, treatment

Vaginal senescence in menopause

Vlad Tica^{1,2,3,4}, Silvia Izvoranu^{1,2,3}, Mihai Banariu^{1,2}, Roxana Penciu^{1,2,3}, Liliana Steriu^{1,2,3}, Corina Nour^{1,2,3}, Diana Mocanu^{1,2,3}, Diana Badiu^{1,2,3}, Irina Tica^{1,3}

1. Faculty of Medicine, "Ovidius" University of Constanța, Romania

2. Institute of Doctoral Studies, Doctoral School of Medicine, "Ovidius" University of Constanța, Romania

3. "Sf. Apostol Andrei" County Emergency Clinical Hospital, Constanța, Romania

4. Academy of Romanian Scientists

Menopause, natural or induced, medically or as a consequence of diseases directly related to the endocrine system, affects the lives of patients, from the physical level to psychoemotional implications. This period has become a challenge for medicine, in its efforts to ensure the health, physical and emotional well-being of women with a growing life expectancy. The paper addresses the effect of hormone deficiency characteristic of the period of perimenopause and menopause on the vagina and the influence on sexuality, with a major impact on the physical and emotional

health of women. The presentation also summarizes the recommendations of the clinical practice guidelines for the control of this specific pathology, as well as the benefits and risks of the recommended therapies, related to the existing or aggravated chronic pathology of menopause. Moreover, psychotherapeutic connotations should be taken into account in the management of the transition from reproductive age to senescence and beyond.

Keywords: menopause, vaginal senescence, management

Is there an endometrium-ovarian cancer link?

Vlad Tica^{1,2,3}, Liliana Steriu^{1,2}, Diana Badiu^{1,2}, Roxana Penciu^{1,2}

1. Faculty of Medicine, Doctoral School of Medicine, "Ovidius" University of Constanța, Romania

2. "Sf. Apostol Andrei" County Emergency Clinical Hospital, Constanța, Romania

3. Academy of Romanian Scientists

Endometriosis is a benign condition defined by the presence of endometrial tissue outside the uterine cavity. This condition is considered a risk factor for ovarian cancer, despite the fact that its presence is a good prognostic factor. The pathology is also a benign estrogen-dependent entity, which has, however, common characteristics with ovarian cancer, such as invasive growth, hormonal dependence and recurrence. The aim of our study was to identify the possible link between endometriosis and associated ovarian cancer. Our research was based on a retrospective analysis of 65 cases of endometriosis hospitalized and operated on for a period of five years, in the Obstetrics-Gynecology I Department of the "Sf. Apostol Andrei" County Emergency Clinical Hospital, Constanța, Romania. Of this group, five patients also had associated ovarian cancer – endometrioid type, high-grade serous type and clear cell carcinoma. These five cases were studied clinically and immunohistochemically (immu-

nomarkers CD10, CD34, Ki67, p53, p16, WT1, estrogen and progesterone receptors). Regarding the low number of cases of endometriosis-associated ovarian cancer, we found no similarities between endometriosis and each type of associated ovarian cancer (low- and high-grade ovarian cancer, respectively). We have observed, on the other hand, elevated levels of estrogen receptors in endometriosis associated with high-grade carcinomas, which may imply a specific role that these receptors play in the carcinogenesis of high-grade carcinomas associated with endometriosis. This analysis can help open the way for patients with endometriosis to identify a risk marker for the development of ovarian cancer, and thus it can lead to the reconsideration of the monitoring of those patients, as well as for targeted therapy.

Keywords: endometriosis, ovarian cancer, immunohistochemistry, CD34, Ki67, p53, estrogen and progesterone receptors, WT1

Correlation between genital prolapse and obesity

Mihaela Camelia Tirnovanu¹, Roxana Mihaela Corduneanu², Andreea Crudu², Vlad Gabriel Tirnovanu³, Bogdan Toma², Alexandra Iov³

1. "Grigore T. Popa" University of Medicine and Pharmacy, Iași, Romania

2. "Cuza Vodă" Clinical Hospital of Obstetrics and Gynecology, Iași, Romania

3. PhD Student, "Grigore T. Popa" University of Medicine and Pharmacy, Iași, Romania

Introduction. Obesity has wide ranging effects on the whole body. The pressure of too much weight can weaken the pelvic floor, resulting in pelvic organs prolapse. **Materials and method.** We made a study on 28 cases with genital prolapse and obesity, admitted in the First Clinic of Gynecology Iași for surgery during the COVID-19 pandemic period. The grade of prolapse and the compartment which descend was correlated with grade of obesity appreciated by Body Mass Index (BMI). We also measured waist circumference, waist-to-hip ratio and the skinfold thickness in three sites – over the triceps in the back of the arm, subscapular and suprailiac. **Results.** The average age was 62.89±8.65 years old (limits: 46-83 years old). Regarding patients with uterine prolapse, 13 presented grade III obesity, 84.6% had grade I obesity and 15.4% had grade II obesity (p=0.734). Ninety percent of the 20 women with cystocele had grade III obesity and 88.9% of them had grade I obesity (p=0.209). Only seven cases presented grade II rectocele and 85.7% had grade I obesity. Twenty-five percent of cases had simultaneously cen-

tral and anterior compartment descent. Three cases were recurrent. As comorbidities caused by obesity, 19 cases had high blood pressure and only seven cases had diabetes mellitus. It should be noted that only 21.4% of patients had ≥4 births. The weight of the baby at delivery was ≥4000 g for 53.3% of the ones with uterine prolapse (p=0.259), 30% of those with cystocele (p=0.067), and 47.6% of those with rectocele (p=0.129). The types of surgery were hysterectomy with bilateral adnexectomy associated with sacropexy in 11 cases or colposuspension in two cases, and colporaphy was performed in 15 cases. **Conclusions.** The demographic characteristics of age, the reproductive status, the number of births and obesity assessment parameters did not differ significantly depending on the type of prolapse. The limit of the present study is represented by the lack of inclusion of patients with incipient degrees of prolapse and obesity, who are treated conservatively.

Keywords: pelvic organ prolapse, obesity, obesity assessment

Clinical and prognostic implications regarding the molecular classification of invasive breast carcinoma

Oana Toader^{1,2}, Alexandra Olaru^{1,2}, Mădălina Mărza¹, Ramona Dragomir^{1,2}, Tiberiu Georgescu^{1,2}

1. Department of Obstetrics and Gynecology, "Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

2. "Polizu" Hospital, "Alessandrescu-Rusescu" National Institute for Mother and Child Health, Bucharest, Romania

Introduction. Invasive breast carcinoma is a heterogeneous group of tumors, including a wide spectrum of morphological variants and molecular alterations with various clinical outcomes in response to different treatment modalities. Although long suspected, prognostic heterogeneity has been molecularly confirmed with the discovery of five intrinsic subtypes. "HER2-low" has been recently described as a novel category of HER2(-) tumors that test IHC 1+ or 2+ and FISH negative, which appear to gain some benefit in clinical trials from HER2-antibody-drug conjugates. **Materials and method.** A literature review has been carried out using the following keywords: "invasive breast cancer", "molecular classification" and "prognosis". Narrative description of the data has been obtained from the selected articles. **Results.** Prognostic gene expression profiles are now commercially available (Oncotype DX, MammaPrint, PAM 50-ROR, EndoPredict, Breast Cancer Index), but the costs, the complexity and the lack of comprehensive data supporting a predictive value for adjuvant therapy hampered their wide adoption in daily practice. Although initially aimed at replacing the standard work-up of immunohistochemi-

cal assessment, it is now increasingly recognized that both methods largely overlap and are most applicable for ER(+), HER2(-) carcinomas. Standard immunohistochemical work-up established by the St. Gallen consensus is cheaper and easier to use and therefore remains the gold standard in estimating the risk for breast cancer relapse and in guiding the clinical decisions regarding efficacy from endocrine, chemotherapy or targeted therapy. **Conclusions.** Studies show that a combination of clinical and pathological features has more prognostic value than each feature alone. Numerous methods of weighting the value of each feature and creating an overall estimate prognosis have been developed, most being relevant for ER(+) cancers: Nottingham Prognostic Index, IHC4 Score, Magee Equations, Flanagan Prognostic Index, Geradts Prognostic Index and Eden Prognostic Index. The early diagnosis and the personalized treatments synergistically contribute to the improvement of prognosis, therefore screening programs maintain a significant role in the era of precision medicine.

Keywords: molecular classification, invasive breast carcinoma, prognosis

Current concepts and etiopathogenic considerations in the diagnosis of pelvic serous tumors

Oana Toader^{1,2}, Ramona Dragomir^{1,2}, Mădălina Mărza¹, Ramona Mircea-Vicol^{1,2}, Tiberiu Georgescu^{1,2}

1. Department of Obstetrics and Gynecology, "Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

2. "Polizu" Hospital, "Alessandrescu-Rusescu" National Institute for Mother and Child Health, Bucharest, Romania

Introduction. High-grade serous carcinoma is one of the most common types of ovarian carcinoma. Based on prevalence and mortality, serous carcinoma is the most important type of ovarian cancer, representing most of the primitive ovarian carcinomas with a negative prognosis. New hypothesis regarding the pathogenesis of ovarian cancer has recently been presented which helps to explain the problems in understanding the development and diverse morphology of these neoplasms. These findings are based on clinicopathologic and molecular genetic studies. **Materials and method.** A review of the literature on the current concepts and etiopathogenic considerations in the diagnosis of pelvic serous tumors has been carried out. The strategy involved the use of keywords such as "ovarian cancer", "low-grade serous carcinoma", "high-grade serous carcinomas" and "molecular pathogenesis", with the selection of articles and the narrative description of the data obtained. **Results.** The development of a dualist model based on clinicopathologic and molecular genetic studies divides all histologic types of epithelial

ovarian carcinomas into two categories: type I and type II. Low grade serous tumors – type I – are most often indolent, diagnosed in stage I, they develop from serous borderline tumors and are genetically stable, being characterized by specific mutations such as KRAS, BRAF and ERBB2, and rarely TP53. High grade serous carcinomas – type II – are diagnosed in advanced stages, are aggressive and develop from intraepithelial carcinomas of the fallopian tube. They have a very high incidence of TP53 mutation and demonstrate marked chromosomal aberrations that remain relatively stable throughout the disease. **Conclusions.** The new hypothesis for the pathogenesis of ovarian serous cancer based on a dualist model and the idea that most of them originate outside the ovary help in organizing the epithelial ovarian neoplasms and lead to the development of new and improved models in preventions, screening and treatment of this affliction.

Keywords: ovarian cancer, low grade serous carcinoma, high grade serous carcinomas, molecular pathogenesis

Multigene NGS panels and cancer susceptibility syndromes – a study report

Radu-Ioan Ursu^{1,2,3}, Diana Prepelitã⁴, Georgiana Ilie⁴, Cristina Dragomir¹, Iuliana Chelu¹, Grația Chelu¹, Simona Filimon¹, Mihaela Popa¹, Bianca Basangiu¹, Andreea Ionescu¹, Andra Perioc¹, Ioana Patricia Riurean¹, Antoanela Curici^{1,4}

1. Central Reference Laboratory, Department of Medical Genetics, Synevo, Romania

2. Department of Medical Genetics, "Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

3. Fetal Medicine Excellence Research Center, "Alessandrescu-Rusescu" National Institute for Mother and Child Health, Bucharest, Romania

4. "Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

Introduction. Since the beginning of the 1990s, with the discovery of the first "cancer genes", numerous genetic factors have been identified through a great number of studies to be associated with the susceptibility for hereditary cancers. Hundreds of genes have been detected, with thousands of variants involved in the development of the disease. Next Generation Sequencing (NGS) technology has been the most important tool in cancer research and cancer diagnosis as well, during the past decades. Most of the laboratories worldwide use multigene NGS panels to identify the genetic causes or the genetic predisposition for cancer. **Materials and method.** NGS sequencing on an Illumina platform has been performed using multigene panels for the different types of cancer. The study included individuals tested during February-September 2021. **Results.** Apart from the BRCA1 and BRCA2 genes mutations, identified in the greater majority of cases, pathogenic variants in other genes, such

as PALB2, CHEK2, MSH6, MLH1, ATM, NF1 etc., have been detected. The most frequent pathology identified in our cohort was breast cancer, followed by colorectal and ovarian cancers. Other types of cancer reported in these individuals were melanoma, cerebral tumors, pancreatic cancer, endocrine tumors, renal cancer, gastrointestinal tumors and prostate cancer. **Conclusions.** NGS panels are the test of choice for the diagnosis of the disease in hereditary cancer patients and for identifying the risk for developing the disease in healthy individuals. Both pre-testing and post-testing genetic counseling are of great importance for the selection of patients for genetic testing and for choosing the most eloquent genetic test for each individual in particular, and also for the clinical interpretation of the results, which in many cases may raise numerous problems and discussions.

Keywords: multigene NGS panels, cancer susceptibility

Particularities of acute pyelonephritis in pregnant women

Cristian Vasilachi

Department of Obstetrics and Gynecology, "Nicolae Testemițanu" State University of Medicine and Pharmacy, Chișinău, Republic of Moldova

Introduction. Acute pyelonephritis (APN) is an inflammatory disease that affects the urinary tract. About 12% of pregnant women suffer from it. APN has as complications septic shock, respiratory insufficiency and kidney failure. **Objective.** To evaluate all functional and morphological changes in pregnant women that can change the ordinary path of the evolution of pyelonephritis, and the impact on pregnancy. **Materials and method.** A retrospective study included data from 45 medical files of pregnant women who were diagnosed with acute pyelonephritis: age (A), number of pregnancies (P), pregnancy trimester (PT), type of lumbar ache (DL), hydronephrosis (H), urinalysis (AGU), results of USG, urine culture (U), and complications – imminent abortion (IA), premature birth (NP), retarded fetal development (RDIU). **Results.** A: 26.5 years old

($p=0.203$), P: 66.7% primiparous ($p=0.278$), PT: 57.8% third trimester ($p=0.310$), DL: 71.1% colicative, on the right side ($p=0.183$), H: 53.3% of cases ($p=0.254$), AGU: leukocyte 100% ($p<0.001$)/RBC – 60.5% ($p=0.130$), bacteria – 57.1% ($p=0.197$), USG: 62% pyelocaliceal dilation ($p=0.174$), U: 95% *E. coli* ($p<0.001$), IA: 40% ($p=0.165$), NP: 15.5% ($p=0.212$), RDIU 13.3% ($p=0.262$); Chi squared test = 29.7. **Conclusions.** Acute pyelonephritis affects pregnant women due to many morphological and functional changes that appear during pregnancy, has a bad influence on the intrauterine development of the child, and elevates the risk of pregnancy disruption, premature delivery, IUD retardation, intrauterine infection development etc.

Keywords: acute pyelonephritis, hydronephrosis, IUD retardation

Treatment of chronic endometritis in cases with implantation failure after *in vitro* fertilization procedure

Cristian Văduva¹, Laurențiu Diră², Sidonia Săndulescu¹, Alice Văduva²

1. University of Medicine and Pharmacy of Craiova, Romania

2. HitMed Medical Center, Craiova, Romania

Introduction. One of the causes of infertility is the inability of the embryo to implant. Chronic endometritis is caused by the action of pathogens that populate the lower genital tract and is a major cause that affects the implantation of the embryo. Several current studies have shown improved fertility after the diagnosis and treatment of endometritis. The present study looked at the effects of endometritis treatment on the rate of *in vitro* fertilization (IVF) pregnancies in couples who had at least one failure in previous procedures. **Materials and method.** The study is a prospective one conducted between May 2018 and May 2021 and was performed on 36 couples with implantation failure after IVF procedures. All these couples underwent control hysteroscopes before and after antibiotic treatment. The hysteroscopic visual aspects, the results of the endometrial biopsies and the sampled intracavitary

cultures were compared. **Results.** Of the 32 cases studied, 19 cases (60.8%) were diagnosed with chronic endometritis either by direct observation or following the histopathological or microbiological result. Antibiotic therapy was a combination of fluoroquinolones and nitroimidazole derivatives. The treatment was performed in repeated cures for three months. In control hysteroscopy, in only two cases (6.2%) no complete cure of endometritis was obtained. Twenty-six cases followed a new IVF procedure, with a positive result in 15 cases (57.7%). **Conclusions.** Hysteroscopic examination of the uterine cavity is particularly important for the success of the IVF procedure. The diagnosis and treatment of chronic endometritis clearly bring benefits to cases that have failed the IVF procedures.

Keywords: endometritis, IVF, implantation, hysteroscopy

Progesterone-follicle ratio is better correlated than blood progesterone level with *in vitro* fertilization results

Cristian Văduva¹, Sidonia Săndulescu¹, Laurențiu Diră², Alice Văduva²

1. University of Medicine and Pharmacy of Craiova, Romania

2. HitMed Medical Center, Craiova, Romania

Introduction. In *in vitro* fertilization (IVF) cycles, a late follicular phase elevation of progesterone (P) is traditionally associated with premature luteinization of endometrium and with a reduced rate of pregnancy. The aim of this study was to investigate the impact of late follicular phase P elevation in relation to ovarian response on cycle outcome. We started from the idea that the average value of P secretion relative to each mature follicle, progesterone-follicle ratio (PFR), better characterizes oocyte fertilization and embryo implantation compared to the value of P alone. **Materials and method.** Data were retrieved from 220 IVF cycles with normal responders in the period 2017-2021. We selected a heterogeneous population undergoing IVF stimulations with a long agonist protocol resulting in

3-15 follicles ≥ 14 mm and a blood P ≤ 3 ng/ml on hCG day followed by fresh embryo transfers. The PFR was calculated for each cycle included by dividing the blood P by the number of follicles ≥ 14 mm. The clinical pregnancy rate was calculated against the range of PFR values and blood P levels. Finally, we associate values of blood P and PFR with clinical pregnancy rate. **Results.** Elevated P levels were associated with a lower pregnancy rate only when they reached $>91^{\text{th}}$ percentile. The PFR was inversely and linearly related to the pregnancy rate for the whole range of values. **Conclusions.** A late follicular increase in blood P is detrimental only if it results from an increase in the average P secreted from each follicle. The limitations are related to study design.

Keywords: progesterone, IVF, follicle

The outcome of structural heart defects diagnosed in the first trimester of pregnancy

Alina Veduță

"Filantropia" Clinical Hospital of Obstetrics and Gynecology, Bucharest

Introduction. We present the results of a detailed protocol of fetal heart examination in the first trimester, in a fetal medicine unit in Bucharest, Romania. **Materials and method.** Since October 2009, in the "Filantropia" Clinical Hospital of Obstetrics and Gynecology, Bucharest, we have systematically assessed pregnancies at 11-14 weeks to screen for aneuploidies and for major fetal structural defects. The fetal anatomy examination protocol included the detailed assessment of the fetal heart. This was performed using the same principles as for the second-trimester examination, in the entire cohort. **Results.** Our population consisted of 7693 patients and 7816 embryos. The protocol for the ultrasound evaluation of the fetal heart was completed for 7597 embryos (97.2%). The outcome is known for 6912 cases (90.9%). We diagnosed 39 heart defects – 30

in the first trimester, seven in the second trimester and two postnatally. Twenty of the 39 heart defects were isolated cardiac malformations. Twelve of the isolated heart defects were diagnosed in the first trimester. The sensitivity of the first-trimester ultrasound in identifying major heart defects was 76.92%. The overall survival in cases of isolated congenital heart disease diagnosed in the first trimester was significantly lower than the survival in the cases diagnosed in the second trimester. **Conclusions.** Many (76.92%) of the significant heart defects can be diagnosed by ultrasound examination, in the first trimester. Our study is an argument for developing a multidisciplinary approach needed for the management of early detected structural heart disease.

Keywords: heart defects, first trimester, ultrasound, protocol, outcome

Management of unexpected *placenta accreta* spectrum after vaginal delivery – case report

Octavia Velicu^{1,3}, Cristina Popescu², Eugen Tănase²

1. Medical Center for Outpatient Diagnosis and Treatment, Ministry of Justice, Bucharest, Romania

2. CF 2 Clinical Hospital, Bucharest, Romania

3. "Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

Placenta accreta spectrum refers to the range of placental pathologic adherence, including *placenta accrete*, *placenta increta* and *placenta percreta*. It affects around 1-200/500 pregnancies, with a rise in numbers seen during the past two decades. The primary diagnostic modality is antenatal diagnosis by obstetric ultrasonography, which leads to planned birth by cesarean in a highly specialized center. The most generally accepted approach to *placenta accreta* spectrum is caesarean hysterectomy with the placenta left *in situ* after

the delivery of the fetus as, usually, attempts at placental removal are associated with a significant risk of hemorrhage. Conservative or expectant management should be rare and considered individually. This paper presents the conservative management of a 33-year-old patient, IIG IP, who delivered vaginally at 36.3 gestational weeks and was afterwards diagnosed with an unexpected *placenta accreta* spectrum.

Keywords: *placenta accreta* spectrum, vaginal delivery, conservative management

The new Enzian classification and the correlation between preoperative evaluation of deep infiltrating endometriosis and intraoperative findings

Simedrea Voicu

EndoInstitute, Premiere Hospital, Timișoara

Introduction. Deep infiltrating endometriosis (DIE) is a debilitating condition with an ill-defined set of symptoms, associated with significant delays in diagnosis and treatment, which severely impacts the patients' quality of life and raises high healthcare costs. **Objective.** The comparison of preoperative stadialization of DIE, by clinical exam, ultrasonography and magnetic resonance imaging (MRI) with intraoperative findings by means of the new Enzian and rASRM score (the revised American Society for Reproductive Medicine). **Materials and method.** Between October 2019 and September 2020, a total of 196 women underwent surgery due to deeply infiltrating endometriosis, at the Premiere Hospital, EndoInstitute – endometriosis competence center specialized in minimally invasive surgery, in Timișoara, Romania. The symptomatology was recorded preoperatively, as well as by clinical exam, intravaginal ultrasonography, MRI, being later compared with the intraoperative findings by means of the ENZIAN score and rASRM score. Laparoscopy was the only surgical approach. **Results.** A clear correlation was seen between the grades of severity in the rASRM score and the Enzian

classification. There was a strong correlation between clinical findings and ultrasonography, in relation with MRI results and intraoperative findings, especially for lesions in vagina, rectovaginal space, rectum, sigmoid colon and bladder, almost 90% for vagina/rectovaginal space and 85% for rectum/sigmoid colon. For uterosacral ligaments and intestine, the sensitivity and negative predictive values were lower, due to technical aspects, such as the impossibility of evaluating the small intestine by ultrasonography. **Conclusions.** In our study, ultrasonography and MRI-based Enzian score correlated well with the intraoperative findings, enabling a better multidisciplinary approach of the surgical procedure, involving endometriosis surgeon, visceral surgeon and urologist. As well, a correct preoperative evaluation of all deep endometriosis lesions is mandatory in order to correctly inform the patients about the possible risks, complications and surgical costs. Therefore, Enzian score represents an excellent communication tool between radiologists and gynecologists, and facilitates the whole process, from diagnosis to surgical treatment.

Keywords: new Enzian classification, endometriosis

Methods of evaluating the colorectal and urinary surgery quality in colorectal and urinary surgery regarding advanced deep infiltrating endometriosis

Simedrea Voicu

EndoInstitute, Premiere Hospital, Timișoara

Introduction. Deep infiltrating endometriosis surgery is a challenging multidisciplinary and complex surgery that requires special skills, harmonized team effort and mastering of anatomy of the patient, along with the physiopathology of the disease. EndoInstitute Timișoara is an excellency European accredited center for surgical treatment of such complex cases. Only in 2020 we have managed surgically 196 DIE cases. Out of these cases, we have performed 172 colorectal surgeries (145 bowel resections, 27 bowel shavings) and 17 ureteral resections (14 end-to-end anastomosis and three ureteral reimplantations). **Materials and method.** We have chosen to evaluate each anastomosis in terms of leakage, vascularization, risk of hemorrhage and tension in tissues. For this purpose, we have used for bowel anastomoses the Michelin test (the classical method of evaluation) and we added the intraoperative endoscopic

inspection for leakage and the ICG test for vascularization. For ureteral anastomosis and replantation, we have assessed the vascularization of the ureteral stomps by ICG test. In some cases, we have used ICF overlay images to determine the ureteral trajectory and nutritive vessels prior to resection step. All ureteral anastomoses were performed on double J catheter that was kept up to 5 weeks after procedure. All the anastomosis (bowel and ureteral) were performed tension-free. We recorded two bowel fistulas and no ureteral fistulas. **Results.** We have concluded that assessing the quality of the anastomoses by using these methods offers a greater safety for the patient and the surgeon, translated in lowering the rates of complications related to anastomotic leakage from 3 to 1.5 percent.

Keywords: colorectal and urinary surgery, anastomosis, infiltrating endometriosis

Low-impact laparoscopy, ERAS and multimodal analgesia in endometriosis advanced surgery

Simedrea Voicu

EndoInstitute, Premiere Hospital, Timisoara

Introduction. Deep infiltrating endometriosis surgery is a challenging multidisciplinary and complex surgery that requires special skills, harmonized team effort and mastering of anatomy of the patient, along with the physiopathology of the disease. EndoInstitute Timișoara is an excellency European accredited center for surgical treatment of such complex cases. Only in 2020, we have managed surgically 196 DIE cases. **Materials and method.** Since the beginning of 2020, we have established special up-to-date protocols for all advanced surgeries, including DIE, in order to minimize the impact of these traumatic and challenging procedures and to maximize the positive experience for the patient and benefit results. These procedures were discussed and agreed in our collective between surgeons (gynecologic, colorectal, urologist and neurosurgeons), pain therapist, anesthesiologist, nutritionist and rehab specialist. All 196 patients were submitted to these procedures that were applied partial or entirely. Regarding low-impact laparoscopy, we have used minilaparoscopy (3.5 mm tro-

cars, 3 mm instruments) and lower pressure CO₂ regime (7-8 mm Hg), along with special dissecting and sealing instruments. Regarding ERAS, we used a cumulus of measures starting with 4-week preoperatively and ending on patient discharge to ensure the best management of the physical and psychological preparation and rehab, bowel preparation and nutrition, fluid, electrolytes and temperature homeostasis, infection prevention and pain management. Regarding multimodal analgesia, we started preoperatively and ended one month after surgery, using local and regional analgesic blocks, non-opioid medication and inhibitors if central pain, along with hypersensitization mechanisms. **Results.** We have evaluated the efficacy of these measures using pain scores and satisfaction questionnaires and we concluded that there is a certain achievement in terms of lowering postoperative pain, enhancing patient satisfaction, and reducing hospitalization and complication rates.

Keywords: ERAS and multimodal, analgesia endometriosis, laparoscopy

ABONARE

Ginecologia.ro



Publicație creditată cu

10 EMC

pentru medici și

5 EMC

pentru moașe,
asistenți maternali și
asistenți medicali



Denumire produs	Detalii produs	Suma	Plată online securizată*
Ginecologia.ro Digital	4 numere format digital	99 lei	
Ginecologia.ro Tipărit	4 numere format tipărit	139 lei	
Combo 1 Digital	2 reviste format digital	165 lei	
Combo 1 Tipărit	2 reviste format tipărit	229 lei	
Combo 2 Digital	Ginecologia.ro + Ziarul Viața Medicală	199 lei	
Combo 2 Tipărit	Ginecologia.ro + Ziarul Viața Medicală	319 lei	


Ordin de plată sau **transfer bancar** pe coordonatele: **MEDICHUB MEDIA SRL**, Green Gate, Bd. Tudor Vladimirescu, nr. 22, etaj 11, sector 5, 050883, București, OP 69 - CP 197, CUI 16136719, J40/2001/2004. Cont IBAN RO73RNCB0617140595120003, deschis la BCR.

*Plată online prin platforma **medichub.ro**

TRATAT *de* PEDIATRIE

EDIȚIA I

Coordonator științific:
PROF. DR. DOINA ANCA PLEȘCA



- 74 DE AUTORI ● 15 SECȚIUNI
- 896 DE PAGINI

Lucrarea se bazează pe experiența dobândită de-a lungul anilor de către autori - în marea lor majoritate, cadre didactice de renume din centre universitare de prestigiu din țară.



400 lei
+ taxă de transport

Pentru comenzi online accesați www.medichub.ro